

## Similarities in the insulating states of twisted bilayer graphene and cuprates

August 1 2019



High resolution scanning tunneling microscope (STM) image of twisted bilayer graphene at the 'magic angle' where electron interactions are maximized. Right: A zoom into the STM image with the corresponding lattices of the twisted bilayer graphene overlayed. Credit: Alexander Kerelsky

In recent decades, enormous research efforts have been expended on the exploration and explanation of high-temperature (high-Tc) superconductors, a class of materials exhibiting zero resistance at



particularly high temperatures. Now a team of scientists from the United States, Germany and Japan explains in *Nature* how the electronic structure in twisted bilayer graphene influences the emergence of the insulating state in these systems, which is the precursor to superconductivity in high-Tc materials.

Finding a material which superconducts at room temperature would lead to a technological revolution, alleviate the <u>energy crisis</u> (as nowadays most energy is lost on the way from generation to usage) and boost computing performance to an entirely new level. However, despite the progress made in understanding these systems, a full theoretical description is still elusive, leaving our search for room temperature superconductivity mainly serendipitous.

In a major scientific breakthrough in 2018, twisted <u>bilayer graphene</u> (TBLG) was shown to exhibit phases of matter akin to those of a certain class of high-Tc superconducting materials—the so-called high-Tc cuprates. This represents a novel inroad via a much cleaner and more controllable experimental setup.

The scientists from the Max Planck Institute for the Structure and Dynamics of Matter (MPSD), Freie Universität Berlin (both in Germany), Columbia University, the Center for Computational Quantum Physics at the Flatiron Institute (both in the U.S.) and the National Institute for Materials Science in Japan focused on the insulating state of TBLG.

This material is made up of two atomically thin layers of graphene, stacked at a very slight angle to each other. In this structure, the insulating state precedes the high-Tc superconducting phase. Hence, a better understanding of this phase and what leads up to it is crucial for the control of TBLG.



The scientists used scanning tunneling microscopy and spectroscopy (STM/STS) to investigate the samples. With this microscopic technique, electrically conducting surfaces can be examined atom by atom. Using the pioneering 'tear and stack' method, they placed two atomically thin layers of graphene on top of one another and rotated them slightly. Then, the team directly mapped the material's atomic-scale structural and electronic properties near the 'magic angle' of around 1.1°.

The findings, which have just been published in *Nature*, cast new light on the factors influencing the emergence of superconductivity in TBLG. The team observed that the insulating state, which precedes the superconducting state, appears at a particular level of filling the system with electrons. This enables scientists to estimate the strength and the nature of the interactions between electrons in these systems—a crucial step towards their description.

In particular, the results show that two distinct van Hove singularities (vHs) in the local density of states appear close to the magic angle which have a doping dependent separation of 40-57 meV. This demonstrates clearly for the first time that the vHs separation is significantly larger than previously thought. Furthermore, the team clearly shows that the vHs splits into two peaks when the system is doped near half Moiré band filling. This doping-dependent splitting is explained by a correlation-induced gap, which means that in TBLG, electron-induced interaction plays a prominent role.

The team found that the ratio of the Coulomb interaction to the bandwidth of each individual vHs is more crucial to the magic angle than the vHs seperation. This suggests that the neighboring superconducting state is driven by a Cooper-like pairing mechanism based on electronelectron interactions. In addition, the STS results indicate some level of electronic nematicity (spontaneous breaking of the rotational symmetry of the underlying lattice), much like what is observed in cuprates near



the superconducting state.

With this research, the team has taken a crucial step towards demonstrating the equivalence of the physics of high-Tc cuprates and those of TBLG materials. The insights gained via TBLG in this study will thus further the understanding of high-temperature superconductivity in cuprates and lead to a better analysis of the detailed workings of these fascinating systems.

The team's work on the nature of the superconducting and insulating states seen in transport will allow researchers to benchmark theories and hopefully ultimately understand TBLG as a stepping stone towards a more complete description of the high-Tc cuprates. In the future, this will pave the way towards a more systematic approach of increasing superconducting temperatures in these and similar systems.

**More information:** Alexander Kerelsky et al. Maximized electron interactions at the magic angle in twisted bilayer graphene, *Nature* (2019). DOI: 10.1038/s41586-019-1431-9

## Provided by Max Planck Institute for the Structure and Dynamics of Matter

Citation: Similarities in the insulating states of twisted bilayer graphene and cuprates (2019, August 1) retrieved 2 May 2024 from <u>https://phys.org/news/2019-08-similarities-insulating-states-bilayer-graphene.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.