

Eight species of fungus cause root rot in South Dakota corn

August 22 2019



The yellowing wilted leaves on these plants are one of the signs of root rot caused by one of eight species of *Fusarium* in South Dakota corn fields, according to a study conducted by researchers in the Department Agronomy, Horticulture and Plant Science. Credit: South Dakota State University

An invisible enemy is attacking South Dakota corn.

The crop may look fine above ground, but as many as eight [species](#) of a

common soil fungus may be infecting the roots—and compromising yields, according to South Dakota State University plant disease experts.

Research associate Paul Okello of the Department of Agronomy, Horticulture and Plant Science identified eight *Fusarium* species that cause root rot in South Dakota cornfields as part of his doctoral research. Furthermore, seven *Fusarium* species isolated from corn also cause disease in soybeans.

This is the first time South Dakota researchers have identified the pathogens causing root rot of corn in the state, according to Field Crops Pathologist Febina Mathew, an assistant professor in the Department of Agronomy, Horticulture and Plant Science. "The number of species can vary from region to region, but eight species have been identified in South Dakota," Okello said.

Fusarium species have a broad host range, which includes soybean, sunflowers and small grains, such as wheat, consequently the pathogen can affect crops that are commonly rotated with corn," said Mathew, who was Okello's research adviser. "Our research on *Fusarium* root rot of corn will help us work with breeders to develop resistant varieties and with chemical companies to test the efficacy of seed treatments that target these specific pathogens."

The May 2019 issue of Plant Health Progress, a journal published by the American Phytopathological Society, featured two articles on the research. The article identifying the *Fusarium* species causing corn root rot received the Editor's Pick Award. Okello was first author on both papers.

The corn research was supported by U.S. Department of Agriculture Hatch Act funding through the South Dakota Agricultural Experiment Station. The soybean work was funded by the South Dakota Soybean

Research and Promotion Council and the North Central Soybean Research Program.

Analyzing corn root rot

While identifying species of *Fusarium* causing root rot of soybean in 2014, Okello noticed previous crops in some of the fields were corn or wheat. That piqued his interest in finding out which *Fusarium* species affected corn.

In 2015, SDSU researchers collected discolored roots from 50 cornfields across a 24-county area in eastern South Dakota, which produces 50 percent of the state's corn.

Estimated yield losses due to corn root rot, seedling blight and [plant-parasitic nematodes](#) during the last six years varied from average of 225,000 bushels per year beginning in 2012 to approximately 75,000 bushels in 2017 and 2018 in South Dakota, according to surveys conducted by the Corn Disease Working Group. A greater awareness of the disease, unfavorable environmental conditions and changes in management strategies may have helped reduce losses recently, Mathew explained.

"*Fusarium* is always in the soil, but environment plays a big role in disease development," Okello said. Cool, wet soil conditions favor disease development. "This year, we are seeing the disease because many of the fields are excessively wet," Mathew added.

Lisa Richardson, executive director of the South Dakota Corn Utilization Council and the South Dakota Corn Growers Association, said, "Though these losses are small relative to the nearly 800 million bushels of corn produced annually in the state, we are thankful that this research helps producers have an awareness and understanding of this

fungus and how it moves and how they can address it should a significant outbreak occur."

Identifying fungal species

Okello isolated the eight species of *Fusarium* from the root specimens and then verified in the greenhouse that these pathogens caused root disease in corn. One of the species, *Fusarium boothii*, had never been reported as affecting corn in the United States. However, *F. boothii* was previously identified as causing disease on wheat in Nebraska.

A 1973 study done by University of Minnesota and Purdue University researchers found six *Fusarium* species in corn. However, the Okello study identified eight species in corn. "This suggests that the *Fusarium* species diversity affecting corn may have changed and additional research is required," he said.

Next, the researchers did a cross-pathogenicity study, putting the *Fusarium* isolates from corn on soybeans and the ones from soybeans on corn in the greenhouse. "We found that seven species of the South Dakota isolates from either soybean or corn cause disease in both crops," Okello said. "This means if you are going to plant soybeans after corn or vice-versa, you are increasing the inoculum level of these soil pathogens, amplifying what's already there."

Mathew's lab is now testing fungicide seed treatment to determine if the current chemistries can help manage *Fusarium* root rot.

"The use of partially resistant hybrids can also decrease losses, Mathew pointed out. "However, these are hybrids that can provide resistance to *Fusarium* ear rot, which growers can verify with the commercial seed companies. At this time, we are not sure if the genes conferring resistance to *Fusarium* [root rot](#) are also responsible for resistance to

Fusarium ear rot and/or stalk rot. This warrants further study."

Though knowing what specific *Fusarium* species infect [corn](#) will help breeders screen germplasm, Mathew said, "the breeding process can be challenging because resistance to *Fusarium* is controlled by multiple genes."

Provided by South Dakota State University

Citation: Eight species of fungus cause root rot in South Dakota corn (2019, August 22) retrieved 1 May 2024 from <https://phys.org/news/2019-08-species-fungus-root-south-dakota.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|