


Where in the universe can you find a black hole nursery?

August 8 2019



Credit: CC0 Public Domain

Gravitational wave researchers at the University of Birmingham have developed a new model that could help astronomers track down the origin of heavy black hole systems in the Universe.

Black holes are formed following the collapse of [stars](#) and possibly supernova explosions. These colossally dense objects are measured in

terms of solar masses (M_{\odot}) - the mass of our sun.

Typically, stars will only form [black holes](#) with masses of up to $45 M_{\odot}$. These systems then pair and merge together, producing gravitational waves that are observed by the LIGO and Virgo detectors.

Stellar collapse, however, causes instabilities that prevent the formation of heavier black holes—so a new model is needed to explain the existence of binary black hole systems with masses larger than about $50 M_{\odot}$.

These objects are thought to be formed from [binary black holes](#) which have then gone on to merge with other black holes. Scientists believe that these 'next generation' black holes —made up of the merger of their 'parents'—might be the heavier black holes observable by LIGO and Virgo.

In a new study, published in *Physical Review D Rapid Communications*, researchers from the University of Birmingham's Institute for Gravitational Wave Astronomy, suggest that future detections of multiple generations of black-hole mergers would allow us to figure out their birthplace. They have produced new calculations that could help astronomers better understand these mergers—and where to find them.

"Star clusters—groups of stars that are bound together by gravity—might act like black-hole 'nurseries', providing an ideal environment to grow generations of black holes, " explains Dr. Davide Gerosa, lead author of the paper. "But in order to know what type of [star clusters](#) are most likely to be capable of producing these, we first need to know something about the physical conditions that would be needed."

The team believes they have found part of the solution to this puzzle by calculating the likely 'escape speed' a [cluster](#) needs to have to be able to

host a black hole with a mass above $50 M_{\odot}$. The escape speed is the velocity at which an object would need to be travelling to escape gravitational pull. For instance, a rocket leaving earth would need to be travelling at 11km/s (25,000 mph) to get into orbit.

When they merge, black holes receive recoils or kick. Much like a gun recoils as a bullet is shot, black holes recoil as [gravitational waves](#) are emitted. The next generation of black holes can form only if their parents have not been 'kicked out' of the cluster, i.e. only if the escape speed of the cluster is large enough.

The team calculated that observing black holes with mass above $50 M_{\odot}$ would suggest that the cluster where they lived had an escape speed larger than about 50km/s.

Co-author Professor Emanuele Berti from Johns Hopkins University, explains: "Gravitational wave observations provide an unprecedented opportunity to understand the astrophysical settings where black holes form and evolve. A very massive event would point towards a dense environment with large escape speed".

Where might you find these types of dense clusters? Many predictions for LIGO and Virgo so far concentrated on 'globular clusters' - spherical collections of about a million stars tightly bound together in the outskirts of galaxies. Their escape speed, however, is too low. This new study finds globular clusters are unlikely to host multiple generations of black holes. Astronomers will need to look further afield: nuclear star clusters, found towards the centre of some galaxies are dense enough and might provide the type of environment needed to produce these objects.

"Gravitational-wave astronomy is revolutionizing our understanding of the Universe," says Dr. Gerosa. "We are all waiting for upcoming results from LIGO and Virgo to put these and other astrophysical predictions to

the test".

More information: arxiv.org/abs/1906.05295

Provided by University of Birmingham

Citation: Where in the universe can you find a black hole nursery? (2019, August 8) retrieved 19 April 2024 from <https://phys.org/news/2019-08-universe-black-hole-nursery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.