


Researchers develop new framework for nanoantenna light absorption

September 23 2019



Qinglan and Cunningham found a way to use nanoantenna technology to harvest light. Credit: Micro and Nanotechnology Laboratory, University of Illinois at Urbana-Champaign

Harnessing light's energy into nanoscale volumes requires novel engineering approaches to overcome a fundamental barrier known as the "diffraction limit." However, University of Illinois researchers have breached this barrier by developing nanoantennas that pack the energy captured from light sources, such as LEDs, into particles with nanometer-scale diameters, making it possible to detect individual biomolecules, catalyze chemical reactions, and generate photons with desirable properties for quantum computing.

The results, which have a broad array of applications that may include better cancer diagnostic tools, were recently published in the *Nano Letters*, a prestigious peer-reviewed journal published by the American Chemical Society in a paper entitled "Microcavity-Mediated Spectrally Tunable Amplification of Absorption in Plasmonic Nanoantennas," The research was funded by the National Science Foundation.

To create a device capable of overcoming the [diffraction limit](#), graduate student Qinglan Huang and her adviser, Holonyak Lab Director Brian T. Cunningham, a Donald Biggar Willett Professor in Engineering, coupled photonic crystals with a plasmonic [nanoantenna](#), an innovative approach in the field. The photonic crystals serve as light receivers and focus the energy into an [electromagnetic field](#) that is hundreds of times greater than that received from the original light source, such as an LED or laser. The nanoantennas, when "tuned" to the same wavelength, absorb the energy from the electromagnetic field and concentrate the energy into a smaller volume that is yet another two orders of magnitude of greater intensity. The energy feedback between the photonic crystal and

the nanoantenna, called "resonant hybrid coupling" can be observed by its effects on the reflected and transmitted light spectrum.

"To get cooperative coupling between two things is exciting because it's never been done," said Huang. "It's a general-purpose concept that we have experimentally demonstrated for the first time."

To achieve this, the team carefully controlled the density of the nanoantennas to maximize their energy collection efficiency. They also developed a method that allowed the nanoantennas to be distributed uniformly across the photonic crystal surface and tuned the [photonic crystal](#)'s optical resonating wavelength to match the absorption wavelength of the nanoantennas.

In addition to changing how researchers can work with light, this new coupling method has the potential to change how and when cancer is diagnosed. One application is to use a gold nanoparticle, not much larger than biomolecules such as DNA, as the nanoantenna. In this case, the feedback provides a way to identify a biomarker unique to a certain type of cancer cell, and the group now linking the resonant hybrid coupling technique to novel biochemistry methods to detect cancer-specific RNA and DNA molecules with single-molecule precision. Cunningham, and other members of the Nanosensor Group will soon publish another paper that focuses specifically on the discovery's applications in regards to cancer diagnostics.

"*Nano Letters* is a very tough journal to get into," said Cunningham. "But the novel physics in this research and the potential for broad applications are what make this research stand out. The next steps of this research involve delving into the potential applications of this new process."

More information: Qinglan Huang et al, Microcavity-Mediated Spectrally Tunable Amplification of Absorption in Plasmonic

Nanoantennas, *Nano Letters* (2019). [DOI: 10.1021/acs.nanolett.9b01764](https://doi.org/10.1021/acs.nanolett.9b01764)

Provided by University of Illinois at Urbana-Champaign

Citation: Researchers develop new framework for nanoantenna light absorption (2019, September 23) retrieved 13 March 2024 from <https://phys.org/news/2019-09-framework-nanoantenna-absorption.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--