

## The first line of defense for Hurricane Dorian is the shoreline

September 4 2019, by Roberto Molar Candanosa



This satellite image, taken on Tuesday, Sept. 3, 2019 and provided by National Oceanic and Atmospheric Administration, shows Hurricane Dorian moving off the east coast of Florida in the Atlantic Ocean. Credit: NOAA

Hurricane Dorian was downgraded to a still-dangerous Category 2



hurricane packing powerful winds and generating catastrophic storm surges that slammed into the Bahamas. After Dorian hit the Abacos Islands on Monday as a Category 5 hurricane, its outer band reached Florida's coast Tuesday evening, and the <u>National Hurricane Center</u> <u>reported</u> that the storm is expected to track north along the East Coast on Wednesday.

Because of the many variables that go into computer modeling for hurricane forecasts, the exact path of a hurricane is <u>extremely difficult</u> to predict. But even if Dorian does not make landfall in the U.S., the <u>storm</u> will create dangerous conditions along the coast. The National Hurricane Center expects <u>up to seven feet of storm flooding</u> in coastal Florida, Georgia, and South Carolina.

As Florida braced for the storm using engineered shorelines to protect from flooding, what may play a key role in buffering some of the <u>flood</u> <u>damage</u> for <u>coastal residents</u> are natural shoreline barriers such as marshes, mangroves, and dunes, says Steven Scyphers, an assistant professor of marine and environmental sciences at Northeastern.

"With our work in the Florida Keys after <u>Hurricane Irma</u>, one thing we found was that coastal residents who had shorelines with natural mangroves typically experienced similar or less damage than folks who would transform the shorelines with bulkheads or seawalls," says Scyphers, who <u>studies people's interactions with the environment along coastlines</u>. "In places where <u>natural habitats</u> can still persist, it's a really important part of a comprehensive strategy for reducing risk."

The National Hurricane Center on Monday evening issued hurricane, storm surge, and tropical storm watches and warnings for the coasts of Florida, Georgia, and South Carolina. Scyphers says one of the main concerns as climate change fuels <u>more intense</u> hurricanes is that substantial segments of U.S. shores that were naturally vegetated with



marshes or mangroves have been lost to residential development.

"There's a really important role in maintaining nature and restoring ecosystems for this risk that we have along the coast," he says. "The shoreline is a small piece of the impact from a storm, but it can also be one that has a lot of connections."

Habitats such as mangroves are well equipped to act as barriers and survive these types of intense storms, often restoring themselves after severe events. That's important for buffering storm surges, which are sometimes considered the deadliest danger from hurricanes for coastal residents.

"Mangroves can be 15 to 16 feet high, and they are basically very dense, almost forest-like barriers that get in the way of the flow of energy and wind," Scyphers says.

Although these natural barriers won't stop a hurricane's devastating force, they play a crucial role in reducing risk for communities where hurricanes are common. Natural barriers are now more than ever an important topic of discussion between scientists and policymakers who work on coastal resilience, Scyphers says.

"That's not the way we've always done it, as we've traditionally relied on building hard infrastructure, which some places, like cities, will always need," he says. "Considering the vulnerability of coastal communities to storms, it's really important to look back at what we are doing to identify what works and what we can do differently."

Many climate models predict <u>more intense</u> and <u>more-rapidly</u> intensifying hurricanes over the next 100 years because of the increasing temperature of the Atlantic. This ability to intensify rapidly is something <u>we have</u> <u>seen</u> in recent years, and 2019 marks the fourth consecutive year that a



Category 5 hurricane has formed in the Atlantic.

It's definitely a concern for people like Katherine Castagno, a joint Northeastern and Nature Conservancy postdoctoral fellow who studies how hurricanes have affected coastal systems over the past 2,000 years. Forecasters say it's <u>the first time a Category 5 hurricane has been</u> <u>generated in four consecutive years</u> in the Atlantic since satellites started tracking storms in the 1960s.

"If you have warmer and warmer water temperatures over where your hurricanes are more likely to form, then you have more energy for those storms to be generated and travel," says Castagno, a postdoctoral fellow in the Social-Ecological Sustainability Laboratory, which is led by Scyphers. "We have seen this in the 2017 Atlantic hurricane season with Irma and Maria, where we have huge storms that are much more intense than what we're used to."

Dorian's winds devastated infrastructure in the Bahamas, and storm surges rose as high as 18 feet. On Tuesday, Bahamas Prime Minister Hubert Minnis <u>confirmed seven deaths</u> on the Abaco Islands. At the height of its strength, the storm was the second-most powerful Atlantic hurricane on record, and is expected to follow a path that will expose a large geographic area to <u>hurricane</u>-force winds along the East Coast.

"It's scary to think that the tragic impacts of these storms will continue as we see more and more of these intense storms," Castagno says.

Provided by Northeastern University

Citation: The first line of defense for Hurricane Dorian is the shoreline (2019, September 4) retrieved 17 May 2024 from <u>https://phys.org/news/2019-09-line-defense-hurricane-dorian-shoreline.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.