


Mother cells as organelle donors

September 13 2019



The Picture depicts the impressive F-actin network (yellow) that is required for the recycling of micronemes (green). It shows an intracellular parasitophorous vacuole in the 8-cell stage that is connected via the “Residual Body”, which is organised by F-actin dynamics. Credit: Dr. Javier Periz, University of Glasgow

Microbiologists at LMU and UoG have discovered a recycling process in the eukaryotic parasite *Toxoplasma gondii* that plays a vital role in the organism's unusual mode of reproduction.

Toxoplasma gondii, the unicellular causative agent of toxoplasmosis, reproduces itself in an unusual fashion by means of an internal budding

process. This entails the development of two [daughter cells](#) within the cytoplasm of the mother cell. On completion of this process, the mother cell undergoes lysis, and the daughter cells are released into the infected host cell. The daughter cells continue to proliferate until the [host-cell](#) itself finally bursts. *T. gondii* is a globally distributed infectious agent. As a rule, the infection is innocuous. However, during pregnancy, transmission of the parasite to the fetus can severely damage the development of the latter. A group of researchers led by Markus Meissner, Professor of Experimental Parasitology at LMU in collaboration with Dr. Javier Periz at Glasgow University, has now described a phenomenon which plays an important role in asexual reproduction—during internal budding, components of a specific organelle are donated by the mother cell to the [daughters](#). The study appears in the online journal *Nature Communications*.

In order to recognize, adhere to and infect host cells, *T. gondii* makes use of organelles called rhoptries and micronemes, which secrete a set of specialized proteins that enable the parasite to invade the target cell. Once the infection has been successfully established, the parasite divides. It had been assumed up to now that the micronemes in the daughter cells are reformed from scratch. However, by specifically labelling one of the micronemal proteins, the authors of the new study were able to follow the fate of the microneme during the cell cycle with the aid of high-resolution microscopy. The observations revealed that the components of the mother cell's microneme are divided more or less equally between the daughter [cells](#). In addition, micronemal proteins are newly synthesized in the daughter cell. The researchers assume that this recycling is not limited to the micronemes, but serves as a more general mechanism to enable the reassembly of organelles that are vital for propagation of the parasite.

"Furthermore, we have shown that recycled micronemes are transported from mother to daughter by the actin filaments of the cytoskeleton," says

Markus Meissner. "This is an entirely new function for actin in the parasite. Up to now, actin was thought to be involved solely in cell motility in *T. gondii*. When we have a better understanding of how this newly discovered function of actin is regulated, we may also be able to identify novel drug targets. This is a very interesting prospect because *T. gondii* is known to possess very few actin-regulating proteins."

More information: Javier Periz et al. A highly dynamic F-actin network regulates transport and recycling of micronemes in *Toxoplasma gondii* vacuoles, *Nature Communications* (2019). [DOI: 10.1038/s41467-019-12136-2](https://doi.org/10.1038/s41467-019-12136-2)

Provided by Ludwig Maximilian University of Munich

Citation: Mother cells as organelle donors (2019, September 13) retrieved 23 April 2024 from <https://phys.org/news/2019-09-mother-cells-organelle-donors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--