


Remora-inspired suction disk mimics fish's adhesion ability, offers evolutionary insight

September 4 2019



Credit: New Jersey Institute of Technology

Remora fishes are famed hitchhikers of the marine world, possessing high-powered suction disks on the back of their head for attaching themselves in torpedo-like fashion to larger hosts that can provide food and safety—from whales and sharks to boats and divers.

Key to the [remora](#)'s adhesion are the disk's well-known capabilities for

generating suction, as well as friction created by spiky bones within the disk called lamellae to maintain hold on its host. However, the factors driving the evolution of remora's unique disc morphology have long eluded researchers seeking to understand, and even engineer new devices and adhesives that mimic, the fish's uncanny ability to lock on to various surface types without harming their host or expending much energy, often for hours at a time under extreme oceanic forces.

In a study led at New Jersey Institute of Technology (NJIT), researchers have showcased a new biologically inspired remora disc capable of replicating the passive forces of suction and friction that power the fish's ability, demonstrating up to 60% greater hold than has been measured for live remoras attached to shark skin.

Using the disc model to explore evolutionary drivers of the remora's disc, researchers say the study's findings provide evidence that today's living species of remora have evolved a greater number of lamellae over time to enhance their holding power and ability to attach to a broader range of hosts with smoother surfaces, thereby increasing their chance for survival.



Opisthomyzon glaronensis fossil featuring six lamellae (top) and dorsal view of a modern remora disc with more rows of lamellae (bottom). According to Brooke Flammang, professor of biological sciences at NJIT, while scientists have shed some light on the origins of the remora's modified fin structure, fundamental aspects of the disc's evolution have largely remained unclear. Credit: Matt Friedman, University of Michigan and Brooke Flammang, New Jersey Institute of Technology

The study, featured in *Bioinspiration and Biomimetics*, indicates the disc model may be used to inform the design of more effective, lower-cost adhesive technologies in the future.

"The beauty behind the remora's adhesive mechanism is that biological tissues inherently do most of the work," said Brooke Flammang, professor of biological sciences at NJIT who led the study. "The most significant aspect of this research is that our robotic disc relies

completely on the [fundamental physics](#) driving the adhesive mechanism in remoras, allowing us to determine biologically relevant performance and gain insight into the evolution of the remora's disc. This was previously not possible with past designs that required a human operator to control the system."

Diverging from many of their closest scavenger-like ancestors, such as cobia (*Rachycentron canadum*), the remora fish (of the family Echeneidae) is believed to have first begun attaching to hosts with rough surfaces, akin to sharks, after having evolved its suction disc from dorsal fin spines nearly 32 million years ago. The disc of living remoras today now features a fleshy-soft outer lip for suction while the disc's interior houses many more linear rows of tissue (lamellae) with tooth-like tissue projections (spinules), which the fish raises to generate friction against various host bodies to prevent slipping during hitchhiking.

According to Flammang, while scientists have shed some light on the origins of the remora's modified fin structure, fundamental aspects of the disk's evolution have largely remained unclear.

"The evolution of the remora's disc is largely unknown," said Flammang. "There is one fossil remora, *Opisthomyzon*, in the fossil record that has a disc with fewer lamellae [than today's remoras] without spinules towards the back of the head."

Flammang says this raises two questions: "how" and "why."

"The 'how' is from the dorsal fin, although the intermediate evolutionary stages aren't known," explained Flammang. "If you look at a phylogeny of remoras it shows that those species that are thought to be more derived have more lamellae ... the 'why' has been assumed to be for adhesive performance, but that was never tested before this paper."

To learn more, Kaelyn Gamel, the study's first author and former graduate researcher in the Flammang lab, designed a remora-inspired disc from commercially available 3-D printed materials that could autonomously maintain attachment to various surfaces and be modified by adding and removing lamellae, enabling the team to investigate the performance of increased lamellar number on shear adhesion.



Image of the team's bioinspired remora disc with modifiable lamellae, constructed using resin-based 3-D printed materials. Credit: New Jersey Institute of Technology

"Our disc's capability to add and remove lamellae while acting as a passive system allowed us to change the amount of friction along with the ambient pressure within the disc," said Gamel, now a Ph.D. researcher at the University of Akron. "We were able to compare the difference between no friction, some friction and a lot of friction based on the variation in lamellae number."

In collaboration with Austin Garner, a researcher at the University of Akron, the team conducted pull-off tests with their model disc underwater, experimenting with the model's lamellar number (up to 12 lamellae) to measure the shear force and time it took to pull the disc from silicon molds with surfaces ranging from completely smooth to those exceeding shark skin roughness (350-grit, 180-grit and 100-grit).

Overall, the team found that their disc's adhesive performance was strongly correlated with an increase in the disc's lamellae, observing a "sweet spot" in suction power between nine and 12 lamellae. When modified to 12 lamellae and 294 spinules, the team's disc weighed just 45 grams and withstood 27 N (newton) forces for 50 seconds—almost three times the force that would typically pull a remora from a shark. The tests also revealed a minimum of six lamellae—the number coincidentally found on the 32-million-year-old fossil *Opisthomyzon*—were needed to maintain adhesion.

"What is most striking about these results is that for a given disc shape, there is an optimal range in which the friction and suction phenomena are balanced, and [as their disc size has gotten longer] remoras have

evolved to maintain this [sweet spot](#) of high-performance adhesion," explained Flammang.

The team now says their remora [disk](#) model will be used for future evolutionary studies to learn whether suction or friction predominated attachment in earliest remora ancestors and how evolution of disc shape affects adhesion. The disc may also have engineering applications in everything from medical biosensors and drug delivery devices to geo-sensing tags for ecological studies and tracking marine life. "One of the greatest advantages to our design is that it operates autonomously because it relies only on the physics of the system for operation," said Flammang. "This makes it easily scalable for a multitude of new technologies, both for medical and scientific purposes."

More information: Kaelyn M Gamel et al, Bioinspired remora adhesive disc offers insight into evolution, *Bioinspiration & Biomimetics* (2019). [DOI: 10.1088/1748-3190/ab3895](https://doi.org/10.1088/1748-3190/ab3895)

Provided by New Jersey Institute of Technology

Citation: Remora-inspired suction disk mimics fish's adhesion ability, offers evolutionary insight (2019, September 4) retrieved 19 April 2024 from <https://phys.org/news/2019-09-remora-inspired-suction-disk-mimics-fish.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.