

## Underwater soundscapes reveal differences in marine environments

September 5 2019



Divers installing an autonomous underwater hydropohone. Credit: Samara Haver

Storms, boat traffic, animal noises and more contribute to the



underwater sound environment in the ocean, even in areas considered protected, a new study from Oregon State University shows.

Using underwater acoustic monitors, researchers listened in on Stellwagen Bank National Marine Sanctuary off the coast of Boston; Glacier Bay National Park and Preserve in Alaska; National Park of American Samoa; and Buck Island Reef National Monument in the Virgin Islands.

They found that the ambient sounds varied widely across the sites and were driven by differences in animal vocalization rates, human activity and weather.

The findings demonstrate that <u>sound</u> monitoring is an effective tool for assessing conditions and monitoring changes, said Samara Haver, a doctoral candidate in the College of Agricultural Sciences at OSU and the study's lead author.

"This is a relatively economical way for us to get a ton of information about the environment," said Haver, who studies marine acoustics and works out of the Cooperative Institute for Marine Resources Studies, a partnership between OSU and the National Oceanic and Atmospheric Administration at the Hatfield Marine Science Center in Newport.

"Documenting current and potentially changing conditions in the <u>ocean</u> soundscape can provide important information for managing the ocean environment."

The findings were published recently in the journal *Frontiers in Marine Science*. Co-authors include Robert Dziak, an acoustics scientist with NOAA who holds a courtesy appointment in OSU's College of Earth, Ocean, and Atmospheric Sciences; and other researchers from OSU, NOAA, Cornell University and the National Park Service.





An autonomous underwater hydrophone deployed in the National Park of American Samoa. Credit: Samara Haver



Passive acoustic monitoring is seen as a cost-effective and low-impact method for monitoring the marine environment. The researchers' goal was to test how effective acoustic monitoring would be for long-term assessment of underwater conditions.

"Ocean noise levels have been identified as a potential measure for effectiveness of conservation efforts, but until now comparing sound across different locations has been challenging," Haver said. "Using equipment that was calibrated across all of the sites, we were able to compare the sound environments of these diverse areas in the ocean."

The researchers collected low frequency, passive acoustic recordings from each of the locations between 2014 and 2018. They compared ambient sounds as well as sounds of humpback whales, a species commonly found in all four locations.

The inclusion of the humpback whale sounds—mostly songs associated with mating in the southern waters, and feeding or social calls in the northern waters—gives researchers a way to compare the sounds of biological resources across all the soundscapes, Haver said.

The researchers found that ambient sound levels varied across all four study sites and sound levels were driven by differences in animal vocalization rates, human activity and weather. The highest sound levels were found in Stellwagen Bank during the winter/spring, driven by higher animal sound rates, vessel activity and high wind speeds. The lowest sound levels were found in Glacier Bay in the summer.

"Generally, the Atlantic areas were louder, especially around Stellwagen, than the Pacific sites," Haver said. "That makes sense, as there is generally more man-made sound activity in the Atlantic. There also was a lot of vessel noise in the Caribbean."



The researchers also were able to hear how sound in the ocean changes before, during and after hurricanes and other severe storms; the monitoring equipment captured Hurricanes Maria and Irma in the Virgin Islands and Tropical Cyclone Winston in American Samoa.

Ultimately, the study provides a baseline for these four regions and can be used for comparison over time. Documenting current and potentially changing conditions in the ocean soundscape can provide important information for managing the ocean environment, particularly in and around areas that have been designated as protected, Haver said.

**More information:** Samara M. Haver et al, Comparing the Underwater Soundscapes of Four U.S. National Parks and Marine Sanctuaries, *Frontiers in Marine Science* (2019). DOI: 10.3389/fmars.2019.00500

Provided by Oregon State University

Citation: Underwater soundscapes reveal differences in marine environments (2019, September 5) retrieved 17 May 2024 from <u>https://phys.org/news/2019-09-underwater-soundscapes-reveal-differences-marine.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.