

## Gas 'waterfalls' reveal infant planets around young star

October 16 2019



Artist impression of gas flowing like a waterfall into a protoplanetary disk gap, which is most likely caused by an infant planet. Credit: NRAO/AUI/NSF, S. Dagnello



The birthplaces of planets are disks made out of gas and dust. Astronomers study these so-called protoplanetary disks to understand the processes of planet formation. Beautiful images of disks made with the Atacama Large Millimeter/submillimeter Array (ALMA) how distinct gaps and ring features in dust, which may be caused by infant planets.

To get more certainty that these gaps are actually caused by planets, and to get a more complete view of planet formation, scientists study the gas in the disks in addition to dust. 99 percent of a protoplanetary disk's mass is gas, of which carbon monoxide (CO) gas is the brightest component, emitting at a very distinctive millimeter-wavelength light that ALMA can observe.

Last year, two teams of astronomers demonstrated a new planet-hunting technique using this gas. They measured the velocity of CO gas rotating in the disk around the young star HD 163296. Localized disturbances in the movements of the gas revealed three planet-like patterns in the disk.

In this new study, lead author Richard Teague from the University of Michigan and his team used new high-resolution ALMA data from the Disk Substructures at High Angular Resolution Project (DSHARP) to study the gas's velocity in more detail. "With the high fidelity data from this program, we were able to measure the gas's velocity in three directions instead of just one," said Teague. "For the first time, we measured the motion of the gas rotating around the star, towards or away from the star, and up- or downwards in the disk."

## Unique gas flows

Teague and his colleagues saw the gas moving from the upper layers towards the middle of the disk at three different locations. "What most likely happens is that a planet in orbit around the star pushes the gas and dust aside, opening a gap," Teague explained. "The gas above the gap



then collapses into it like a waterfall, causing a rotational flow of gas in the disk."

This is the best evidence to date that there are indeed planets being formed around HD 163296. But astronomers cannot say with one hundred percent certainty that the gas flows are caused by planets. For example, the star's magnetic field could also cause disturbances in the gas. "Right now, only a direct observation of the planets could rule out the other options. But the patterns of these gas flows are unique and it is very likely that they can only be caused by planets," said co-author Jaehan Bae of the Carnegie Institution for Science, who tested this theory with a computer simulation of the disk.

The location of the three predicted planets in this study correspond to the results from last year: they are likely located at 87, 140 and 237 AU. (An astronomical unit—AU—is the average distance from the Earth to the Sun.) The closest planet to HD 163296 is calculated to be half the mass of Jupiter, the middle planet is Jupiter-mass, and the farthest planet is twice as massive as Jupiter.





Scientists measured the motion of gas (arrows) in a protoplanetary disk in three directions: rotating around the star, towards or away from the star, and up- or downwards in the disk. The insert shows a close-up of where a planet in orbit around the star pushes the gas and dust aside, opening a gap. Credit: NRAO/AUI/NSF, B. Saxton





An artist's conception of the disk of gas and dust rotating around the young star HD 163296. Gas can be seen cascading into gaps in the disk--likely indicating the formation of baby planets in these locations. Credit: Robin Dienel, Carnegie Institution for Science.

## **Planet atmospheres**

Gas flows from the surface towards the midplane of the protoplanetary disk have been predicted by theoretical models to exist since the late '90s, but this is the first time that they have been observed. Not only can they be used to detect infant planets, they also shape our understanding of how gas giant planets obtain their atmospheres.

"Planets form in the middle layer of the disk, the so-called midplane. This is a cold place, shielded from radiation from the star," Teague



explained. "We think that the gaps caused by planets bring in warmer gas from the more chemically active outer layers of the disk, and that this gas will form the atmosphere of the planet."

Teague and his team did not expect that they would be able to see this phenomenon. "The disk around HD 163296 is the brightest and biggest disk we can see with ALMA," said Teague. "But it was a big surprise to actually see these gas flows so clearly. The disks appears to be much more dynamic than we thought."

"This gives us a much more complete picture of planet formation than we ever dreamed," said co-author Ted Bergin of the University of Michigan. "By characterizing these flows we can determine how planets like Jupiter are born and characterize their chemical composition at birth. We might be able to use this to trace the birth location of these planets, as they can move during formation."

**More information:** Meridional flows in the disk around a young star, *Nature*, DOI: 10.1038/s41586-019-1642-0, nature.com/articles/s41586-019-1642-0

Provided by National Radio Astronomy Observatory

Citation: Gas 'waterfalls' reveal infant planets around young star (2019, October 16) retrieved 1 May 2024 from <u>https://phys.org/news/2019-10-cascades-gas-young-star-early.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.