

# Lakes worldwide are experiencing more severe algal blooms

October 14 2019

---



Lake Khanka (left) on the border between Russia and China and Lake St. Clair (right) on the border between the United States and Canada both showed improvement and then deteriorated in Ho, Michalak, and Pahlevan's study of algal blooms in large lakes. Credit: Khanka image made by Norman Kuring, NASA's Ocean Color web, and Lauren Dauphin. St. Clair image courtesy of NASA Earth Observatory made by Joshua Stevens, using Landsat data from the U.S. Geological Survey.

The intensity of summer algal blooms has increased over the past three decades, according to a first-ever global survey of dozens of large, freshwater lakes, which was conducted by Carnegie's Jeff Ho and Anna Michalak and NASA's Nima Pahlevan and published by *Nature*.

Reports of harmful [algal blooms](#)—like the ones that shut down Toledo's water supply in 2014 or led to states of emergency being declared in Florida in 2016 and 2018—are growing. These aquatic phenomena are harmful either because of the intensity of their growth, or because they include populations of toxin-producing phytoplankton. But before this research effort, it was unclear whether the problem was truly getting worse on a global scale. Likewise, the degree to which human activity—including agriculture, urban development, and [climate change](#)—was contributing to this problem was uncertain.

"Toxic algal blooms affect drinking water supplies, agriculture, fishing, recreation, and tourism," explained lead author Ho. "Studies indicate that just in the United States, freshwater blooms result in the loss of \$4 billion each year."

Despite this, studies on freshwater algal blooms have either focused on individual lakes or specific regions, or the period examined was comparatively short. No long-term global studies of freshwater blooms had been undertaken until now.



Ho, Michalak, and Pahlevan's study of algal blooms in lakes over a 30-year period found that Florida's Lake Okeechobee deteriorated. Toxic algal blooms resulted in states of emergency being declared in Florida in 2016 and 2018. Credit: NASA Earth Observatory image made by Joshua Stevens, using Landsat data from the U.S. Geological Survey.

Ho, Michalak, and Pahlevan used 30 years of data from NASA and the

U.S. Geological Survey's Landsat 5 near-Earth satellite, which monitored the planet's surface between 1984 and 2013 at 30 meter resolution, to reveal long-term trends in summer algal blooms in 71 large lakes in 33 countries on six continents. To do so, they created a partnership with Google Earth Engine to process and analyze more than 72 billion [data points](#).

"We found that the peak intensity of summertime algal blooms increased in more than two-thirds of lakes but decreased in a statistically significant way in only six of the lakes," Michalak explained. "This means that algal blooms really are getting more widespread and more intense, and it's not just that we are paying more attention to them now than we were decades ago."



Lago de Cahora Bassa in Mozambique was one of the lakes that Ho, Michalak, and Pahlevan's study of the intensity of lakewater algal blooms showed had sustained improvement over the 30-year period. Credit: NASA Earth Observatory image made by Jesse Allen, using MODIS data from the Land Atmosphere Near-real-time Capability for EOS (LANCE) and Landsat data from the U.S. Geological Survey.

Although the trend towards more-intense blooms was clear, the reasons for this increase seemed to vary from [lake](#) to lake, with no consistent patterns among the lakes where blooms have gotten worse when considering factors such as fertilizer use, rainfall, or temperature. One clear finding, however, is that among the lakes that improved at any point over the 30-year period, only those that experienced the least warming were able to sustain improvements in [bloom](#) conditions. This suggests that climate change is likely already hampering lake recovery in some areas.

"This finding illustrates how important it is to identify the factors that make some lakes more susceptible to climate change," Michalak said. "We need to develop water management strategies that better reflect the ways that local hydrological conditions are affected by a changing climate."

**More information:** Widespread global increase in intense lake phytoplankton blooms since the 1980s, *Nature* (2019). [DOI: 10.1038/s41586-019-1648-7](#) , [nature.com/articles/s41586-019-1648-7](https://www.nature.com/articles/s41586-019-1648-7)

Provided by Carnegie Institution for Science

Citation: Lakes worldwide are experiencing more severe algal blooms (2019, October 14) retrieved 26 April 2024 from <https://phys.org/news/2019-10-lakes-worldwide-experiencing-severe-algal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.