


Magneto-inertial fusion experiment nears completion

October 21 2019




Supersonic jets fired from 7 plasma guns collide in PLX test firings. In addition to debugging the machine, the experiments provide data to verify models of colliding plasmas important for astrophysics, aeronautics, and various controlled fusion schemes. Credit: Los Alamos National Laboratory

Assembly of the Plasma Liner Experiment (PLX) at Los Alamos National Laboratory is well underway with the installation of 18 of 36 plasma guns in an ambitious approach to achieving controlled nuclear fusion (Figure 1). The plasma guns are mounted on a spherical chamber, and fire supersonic jets of ionized gas inward to compress and heat a central gas target that serves as fusion fuel. In the meantime, experiments performed with the currently installed plasma guns are providing fundamental data to create simulations of colliding plasma jets, which are crucial for understanding and developing other controlled fusion schemes.

Most [fusion](#) experiments employ either magnetic confinement, which relies on powerful magnetic fields to contain a fusion [plasma](#), or inertial confinement, which uses heat and compression to create the conditions for fusion.

The PLX machine combines aspects of both magnetic confinement fusion schemes (e.g. tokamaks) and inertial confinement machines like the National Ignition Facility (NIF). The hybrid approach, although less technologically mature than pure magnetic or inertial confinement concepts, may offer a cheaper and less complex fusion reactor development path. Like tokamaks, the fuel plasma is magnetized to help mitigate losses of particles and thermal energy. Like inertial [confinement](#) machines, a heavy imploding shell (the plasma [liner](#)) rapidly compresses and heats the fuel to achieve fusion conditions. Instead of NIF's array of high-power lasers driving a solid capsule, PLX relies on supersonic plasma jets fired from plasma guns.

The PLX has an additional advantage: Because the fusion fuel and liner are initially injected as a gas, and the plasma guns are located relatively far from the imploding fuel, the machine can be fired rapidly without damage to the machine components or the need for replacement of costly machined targets.



Half of the 36 supersonic plasma jets that will make up the Plasma Liner Experiment at Los Alamos National Laboratory have been installed. The remaining jets will be added and fully spherical plasma liner experiments will commence by the end of 2020. Credit: Los Alamos National Laboratory

"We will conduct experiments this year to study the formation of a hemispherical liner with 18 guns installed," said Dr. Samuel Langendorf, a scientist with the lab's Experimental Physics Group who is leading the assembly of PLX. "We hope to complete the installation of the remaining 18 guns in early 2020 and to be conducting fully spherical experiments by the end of 2020. This will allow us to measure the scaling of the liner ram pressure on stagnation as well as the liner uniformity, which are important metrics of the liner performance."

In its partially completed state, the PLX guns are proving useful in studies that Dr. Tom Byvank is performing on colliding plasmas (Figure 2).

"Different models show discrepancies in the simulations of plasma collisions involving multiple ion species," said Dr. Byvank, a postdoc in the Experimental Physics Group. "Our experimental observations of these plasmas help to validate simulations that are important for understanding high-energy-density, supersonic plasmas encountered in astrophysics, aerodynamics and various plasma fusion [machines](#), including the PLX magneto-inertial fusion approach and possibly also [inertial confinement](#) designs like the National Ignition Facility."

More information: Abstracts:

Experimental Observation of Shock-front Separations in Multi-ion-species Collisional Plasma Shocks

2:00 PM - 5:00 PM, Monday, October 21, 2019

Room: Floridian Ballroom AB

meetings.aps.org/Meeting/DPP19/Session/CI2.5

Progress toward the formation of fully spherical imploding plasma liners on PLX

9:30 AM - 12:30 PM, Thursday, October 24, 2019

Room: Grand D

meetings.aps.org/Meeting/DPP19/Session/TO6.12

Provided by American Physical Society

Citation: Magneto-inertial fusion experiment nears completion (2019, October 21) retrieved 9 April 2024 from <https://phys.org/news/2019-10-magneto-inertial-fusion-nears.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.