


New method gives first global picture of mutual predictability of atmosphere and ocean

October 7 2019



Credit: UMD

University of Maryland (UMD) scientists have carried out a novel statistical analysis to determine for the first time a global picture of how the ocean helps predict the low-level atmosphere and vice versa. They observed ubiquitous influence of the ocean on the atmosphere in the extratropics, which has been difficult to demonstrate with dynamic models of atmospheric and oceanic circulation. The results are published today in the *Journal of Climate*, "Local atmosphere-ocean predictability: dynamical origins, lead times, and seasonality."

The research draws on a classic statement often heard in introductory statistics classes that "correlation is not causation." Clive Granger was a Nobel-laureate mathematician who came up with a novel method to address this issue by distinguishing correlation from causation.

"The Granger method relies upon a simple but important notion that a cause precedes its effect, and should improve the prediction of its effect in the future. We realized that this could be a powerful method to study the interactions between [atmosphere](#) and [ocean](#), and to provide a global picture of how well they predict each other," said applied mathematician Safa Motesharrei, an Environmental Systems Scientist at UMD. "This method sheds light on both the potential to better predict regional climate as well as the nature of the interactions."

"There are many physical processes that govern the interaction between the atmosphere and ocean," said lead author Eviatar Bach, Ph.D. student in the Department of Atmospheric and Oceanic Science (AOSC) at UMD. "For example, wind blowing on the [ocean surface](#) creates currents, and the sea surface heats up the lower atmosphere. These interactions between the atmosphere and ocean play a major role in climate and our ability to predict it, so understanding their geographical structure is important."

"It has been known that in the tropical oceans, the ocean is predominantly driving the atmospheric changes, while in the extratropics the atmosphere generally drives the ocean," said co-author Eugenia Kalnay, Distinguished University Professor of AOSC at UMD. "I developed a dynamical rule to determine the direction of the forcing in 1986, and others have addressed this question using climate models. This study provides a definitive answer."

The basic Granger method was introduced in 1969, but the authors "cleverly applied it for the first time to atmosphere and ocean data," said

Juergen Kurths, Head of Complexity Science Department at Potsdam Institute for Climate Impact Research in Germany, who was not a co-author. Kurths is a prominent physicist who has developed many novel mathematical methods for studying climate and other nonlinear systems.

"The most novel finding of this research is that the method of Granger causality found the ocean to influence the atmosphere almost everywhere in the extratropics," said Samantha Wills, a postdoctoral researcher at NOAA's Pacific Marine Environmental Laboratory, who was not a co-author. "This can be a challenging task given that the atmosphere dominates air-sea interaction in the extratropics, and the influence of the ocean on the atmosphere is not much larger than internal variability."

"This had not been demonstrated by previous General Circulation Model experiments. Although there have been a few special cases where it has been shown that mid-latitude sea-surface temperatures have a significant impact on the atmosphere, this relationship was not known to be as ubiquitous as this paper has shown," said J. Shukla, University Professor at George Mason University, who was not a co-author. Shukla is a world renowned climate scientist who pioneered studies of predictability.

Moreover, the study's estimates of the spatial structure of predictability could help to further advance the science of coupled data assimilation, the nascent field that attempts to leverage the interactions between atmosphere and ocean to improve [climate](#) prediction.

"The ability to anticipate changes to the ocean or atmosphere based on information from the other system provides society with the opportunity to prepare for future impacts, such as to agriculture and fisheries," said Wills.

"This is a very important paper in the history of predictability research,"

said Shukla, "It will surely inspire further research by the predictability research community. In particular, this paper identifies geographical regions on the globe over which there exists potential predictability which can be harvested for improving operational predictions."

More information: Eviatar Bach et al, Local Atmosphere–Ocean Predictability: Dynamical Origins, Lead Times, and Seasonality, *Journal of Climate* (2019). [DOI: 10.1175/JCLI-D-18-0817.1](https://doi.org/10.1175/JCLI-D-18-0817.1)

Provided by University of Maryland

Citation: New method gives first global picture of mutual predictability of atmosphere and ocean (2019, October 7) retrieved 17 July 2024 from <https://phys.org/news/2019-10-method-global-picture-mutual-atmosphere.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.