

Researchers develop model for more realistic risk assessment of pesticides

October 30 2019

Toxic substances such as pesticides can cause effects on sensitive individuals in concentrations up to 10,000 times lower than previously assumed. This was shown by Researchers at the Helmholtz Centre for Environmental Research (UFZ) in their latest study published in *Scientific Reports*. For understanding these results, the level of stress plays an important role. Most surprisingly: Either too much or too little stress can lead to a higher sensitivity to toxicants.

Whether a toxicant harms or kills an organism, and to which extent, depends on its concentration and the sensitivity of the individuals exposed: The dose makes the poison.

"But that's not all," says UFZ ecotoxicologist Prof. Dr. Matthias Liess. "Also, the magnitude of environmental <u>stress</u> is very important to determine the effect of a toxicant." Toxicant concentration, individual sensitivity and environmental stress thus interact in a triad of effects.

In their current study, the researchers have investigated the role of these individual components. They wanted to find out how sensitive individuals can be better protected by risk assessment. "Previous model calculations could not sufficiently predict the effects of low toxicant concentrations on sensitive individuals and species," explains Liess. "But this is very important in both human and ecotoxicological risk assessment". This is because sensitive persons—such as children, sick or elderly people—or more sensitive species of an ecosystem obviously suffer damage even at much lower concentrations than previously



assumed.

"Our extensive investigation was triggered by an observation: At very low pesticide concentrations—far below concentrations that led to effects in previous studies—effects on sensitive organisms were observed," says the ecotoxicologist. These effects have not been previously observed because such low concentrations are very rarely tested.

Previously, it was assumed that pollutants only elicit effects in high concentrations after a threshold value has been exceeded. However, at such extremely low concentrations, this is obviously not the case. Liess said, "We observed these unexpected effects at very low concentrations in almost all existing studies in which effects of such concentrations of toxicants were investigated. Also, in their own investigations in which they exposed the crustacean Daphnia magna to very low concentrations of the pesticide esfenvalerate, used as an insecticide with contact and feeding action in plant protection products in fruit, vegetable and arable farming in the E.U. The question that arises is: Which process can induce effects in these low concentration ranges? The hypothesis of the UFZ scientists is that toxicant stress meets internal stress. But what does that mean?

Under the influence of environmental stress such as predation pressure, parasites and heat organisms become more sensitive to toxicants and can therefore be affected, or die at much lower toxicant concentrations than in the absence of environmental stress. "We were already able to quantify this relationship in earlier studies," says Liess. "Additionally, we are now able to show that individuals develop internal stress when they are exposed to too little stress from the environment. In fact, it seems like organisms are adapted to a certain degree of 'external' stress. If it is missing, they develop 'internal' stress.



"And since external and internal stress add up, the sensitivity to toxicants increases drastically," explains Liess. The result: Sensitive individuals react to extremely low concentrations of toxicants. These can be up to 10,000 times lower than the concentrations previously considered harmful so far. "Thus, too much, but also too little stress increases sensitivity to pollutants," says the UFZ researcher. "Optimal with regard to resilience against toxicant exposure would therefore be some fluctuating <u>environmental stress</u>, which reduces internal stress.

In order to make effects visible at low concentrations, the scientists developed a model that enables a calculation of the internal stress and the resulting survival. Liess says, "We hope that our study will contribute to a more realistic environmental and human risk assessment—especially for sensitive individuals".

More information: Matthias Liess et al. Predicting low-concentration effects of pesticides, *Scientific Reports* (2019). DOI: 10.1038/s41598-019-51645-4

The model of the UFZ researchers is available online: <u>www.systemecology.eu/indicate/</u>

Provided by Helmholtz Association of German Research Centres

Citation: Researchers develop model for more realistic risk assessment of pesticides (2019, October 30) retrieved 7 May 2024 from <u>https://phys.org/news/2019-10-realistic-pesticides.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.