


Scientists build a nanocage with antiaromatic walls

October 23 2019



(a) construction of antiaromatic-walled nanospace. (b) X-ray crystal structure with a 3D NICS grid, showing magnetic deshielding experienced within the nanospace. Antiaromaticity effects becomes stronger in the order of yellow

Researchers at Tokyo Institute of Technology, the University of Cambridge, and the University of Copenhagen have built a self-assembled nanocage with a very unusual nanospace: Its walls are made of antiaromatic molecules, which are generally considered too unstable to work with. By overturning assumptions about the limits of nano-chemical engineering, the study creates an entirely new nanospace for scientists to explore. Nanometer-sized cavities are already finding a range of useful applications in chemistry, medicine and environmental science.

Scientists including Masahiro Yamashina of Tokyo Institute of Technology (JSPS Overseas Research Fellow, at that time) and Jonathan R. Nitschke of the

University of Cambridge, reporting their work in the journal *Nature*, describe the construction of a new type of nanospace inside "a self-assembled cage composed of four metal ions with six identical antiaromatic walls."

Until now, many teams have developed nanocages with aromatic walls, but none with antiaromatic [compounds](#), owing to the challenges posed by their inherent instability. Aromaticity refers to a property of ring-shaped [organic compounds](#) that makes them highly stable, whereas antiaromaticity describes compounds that are far more reactive, due to a difference in the number of so-called π -electrons shared by the ring.

The team's search for a suitable building block for their nanocage led them to a 2012 study by Hiroshi Shinokubo and co-workers in Japan. This study reported the synthesis of an unusually stable, nickel-based antiaromatic compound called norcorrole. Then, drawing on Jonathan R. Nitschke and his group's expertise in subcomponent self-assembly, the team succeeded in building a three nanometer-diameter cage with a norcorrole skeleton.

To investigate the degree of antiaromaticity within the cage, the team performed nucleus-independent chemical shift (NICS) calculations. The results indicated that the norcorrole panels appear to work together to enhance antiaromaticity. The NICS value was consistently high in the central part of the cage, suggesting that the panels reinforce each other.

The unique environment inside the cage was further tested by encapsulating a series of guest molecules, beginning with coronene which has been already encapsulated within the aromatic cage.

The researchers hypothesized that when exposed to an [external magnetic field](#), guest molecules in an aromatic-walled cage would experience a shielding effect, while those in an antiaromatic-walled cage would experience a deshielding effect.

As predicted by theory, nuclear magnetic resonance (NMR) spectroscopy analyses revealed a deshielding effect attributable to the antiaromatic walls.

All guest molecules tested in the study showed significant downfield chemical

shifting, an indicator of the degree of deshielding. The shift differences ranged from 0.7 to 14.9 parts per million. Of these, a carbon nanobelt showed the highest degree of downfield shifting observed so far resulting from an antiaromatic environment.

The [cage](#) can be considered as a new type of NMR shift reagent, the researchers say, meaning that it could be a useful tool for structural analysis, ie for interpreting the finest structures of organic compounds.

Future work will focus on investigating chemical reactivity within the nanospace.

More information: An antiaromatic-walled nanospace, *Nature* (2019). [DOI: 10.1038/s41586-019-1661-x](https://doi.org/10.1038/s41586-019-1661-x) , [nature.com/articles/s41586-019-1661-x](https://www.nature.com/articles/s41586-019-1661-x)

Provided by Tokyo Institute of Technology

Citation: Scientists build a nanocage with antiaromatic walls (2019, October 23) retrieved 24 April 2024 from <https://phys.org/news/2019-10-scientists-nanocage-antiaromatic-walls.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.