

# First South American insect that emits blue light is discovered

October 31 2019, by André Julião

---



Larvae of a fungus gnat found in Iporanga, São Paulo State, Brazil, have bioluminescent properties previously observed only in species native to North America, New Zealand and Asia. This study paves the way for new biotech applications. Credit: Henrique Domingos / IPBio

Brazilian researchers have discovered a new species of fungus gnat

(Keroplastidae) whose larvae emit blue light. The small fly inhabits an Atlantic Rainforest reserve in São Paulo State. This is the first record of a blue bioluminescent species in the Neotropics. Many bioluminescent insects and fungi have been studied in the region, but all emit green, yellow or red light. The new species has been named *Neoceroplatus betaryiensis* and is described in an article in *Scientific Reports*.

"The larvae were found while bioluminescent mushrooms were being collected and drew attention because they emitted blue [light](#). Fungi and fireflies don't emit blue light, so it had to be a [new species](#)," Cassius Stevani, a professor at the University of São Paulo's Institute of Chemistry (IQ-USP) and last author of the article, told.

The study was part of the Thematic Project "Electronic chemiexcitation in [biological systems](#): bioluminescence and photochemistry in the dark", for which Etelvino José Henriques Bechara, a professor at IQ-USP, is principal investigator.

According to Stevani, [species](#) that emit blue light had previously been found only in North America, New Zealand and Asia. This one was found in Reserva Betary, a privately held Atlantic Rainforest reserve in Iporanga, São Paulo State, bordering the Upper Ribeira State Tourist Park (PETAR).

Biologists Isaias Santos and Grant Johnson, a US-born technical trainee with a scholarship from São Paulo Research Foundation—FAPESP, participated in the collection expedition. Both work at the Biodiversity Research Institute (IPBio), a nongovernmental organization that runs Reserva Betary, offering tourism, environmental education and research opportunities there. Many of the world's species of bioluminescent mushrooms can be found on the property.

The new species of bioluminescent insect was described by entomologist

Rafaela Falaschi, currently a postdoctoral researcher at the University of Ponta Grossa (UEPG). The species epithet (*betaryiensis*) refers to the Betary, a tributary of the Ribeira.

## **Different light patterns**

According to Stevani, adults of the species do not emit light. The bioluminescent larvae live hidden in tree trunks and have three lanterns, one in the tail and two near the eyes.

However, one of the specimens collected by the researchers emitted light at various points on its body. The larva was taken to the laboratory, where it pupated. The pupa remained bioluminescent, but a wasp emerged instead of a fungus gnat.

The researchers concluded that the wasp also belongs to a new species in the parasitoid family Ichneumonidae, which lay eggs in the larvae of beetles, moths and other insects. It is unclear, however, if the different pattern of light emission observed in the specimen is due to an infection caused by the wasp, if it indicates a new species of gnat, or if the pattern is related to sexual dimorphism in *N. betaryiensis* (different morphological traits in males and females).

## **New system**

In addition to the importance of any new species to the production of knowledge about biodiversity, insects that emit blue light are extremely rare, and the discovery suggests the possibility of unearthing a new system of bioluminescence that could have applications in analysis and biotechnology, such as in marking specific cells or genes in biological studies or pollution biosensors, for example.

Like all bioluminescent organisms, the new species generates light via a reaction between luciferin, a substrate, and luciferase, an enzyme that catalyzes it. In assays performed to study the phenomenon, the two compounds are typically isolated by producing an extract of the insect and separating it into portions. One is stored on ice to preserve its enzymes, including luciferase. The other is heated to eliminate the enzymes and leave only the substrate luciferin.

To begin characterizing the bioluminescent system in *N. betaryiensis*, the research group led by [Vadim Viviani](#), a professor at the Federal University of São Carlos (UFSCar) in Sorocaba, São Paulo State, used luciferin and luciferase purified in their laboratory from another species capable of emitting blue light, *Orfelia fultoni*, which inhabits the Appalachian Mountains in the US and Canada.

"Thanks to having luciferase and luciferin from *O. fultoni* already purified in our lab, we were able to analyze cross-reactions with the new species. Light was emitted in all combinations. We also show that larvae of this fungus gnat contain a luciferin-storing protein known as SBF, which is short for substrate-binding fraction, as does *O. fultoni*. Therefore, both species have the same biochemical system," said Viviani, who leads a research group on bioluminescence and biophotonics funded by the National Council for Scientific and Technological Development (CNPq), a federal government agency.

In 2000, Viviani and researchers Thérèse Wilson and J. Woodland Hastings produced the first characterization of *O. fultoni*'s bioluminescent system while Viviani was a postdoctoral fellow at Harvard University in the US. Since then, he has worked steadily on the biochemical characterization of luciferin and luciferase in these insects.

Viviani is the principal investigator for the Thematic Project "Arthropod bioluminescence" funded by FAPESP. His group recently discovered a

species of the genus *Neoditomyia* in the caves of Intervalles State Park, an Atlantic Rainforest remnant in the south of São Paulo State. This species produces luciferin and its SBF but does not emit light. When the substrate was blended with luciferase from *O. fultoni* and the new species, however, it emitted [blue light](#).

Genetic analysis also shows that the new species is closely related to *Neoditomyia* and *O. fultoni*.

Based on their knowledge of other species, the researchers now plan to isolate and investigate the luciferin and luciferase produced by *N. betaryiensis*, which is rarer and less easily found than the North American species.

"We already have luciferin and luciferase from *O. fultoni*, the North American species, and from the *Neoditomyia* found in Intervalles. They have been purified and partially characterized by our lab at UFSCar. This enabled us to conduct this initial study and will also facilitate the isolation of luciferin and cloning of luciferase from *Neoceroplatus* in the future," Viviani said.

Once isolated, the substances will be cloned, and their structure will be determined. The group at UFSCar will work on the luciferase, and Stevani's group at IQ-USP will be responsible for analyzing the luciferin.

"We already have the molecular formula for luciferin. We know how many atoms of carbon, hydrogen, nitrogen, oxygen, sulfur and other elements it contains, but we don't know how these atoms are linked. We need to perform experiments using nuclear magnetic resonance imaging, among other techniques, in order to elucidate the structure. I expect that to happen soon," said Stevani.

**More information:** Rafaela L. Falaschi et al, *Neoceroplatus*

betaryiensis nov. sp. (Diptera: Keroplatidae) is the first record of a bioluminescent fungus-gnat in South America, *Scientific Reports* (2019).  
[DOI: 10.1038/s41598-019-47753-w](https://doi.org/10.1038/s41598-019-47753-w)

Provided by FAPESP

Citation: First South American insect that emits blue light is discovered (2019, October 31)  
retrieved 24 April 2024 from  
<https://phys.org/news/2019-10-south-american-insect-emits-blue.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.