

Research indicates abundance of microbe diversity key to healthy coastal ecosystem

November 5 2019, by Laura Schmitt



Small mixed lucinid clams collected from Sugarloaf Key, Florida, in 2016.
Credit: Annette Engel, University of Tennessee, Knoxville

For millions of years, symbiotic bacteria have lived inside the gill cells of *Lucinidae* clams found in seagrass meadows located mainly along

tropical coasts, such as the Florida Keys. These bacteria play a crucial role in the clam's survival while also contributing to the overall health of the seagrass in which the clams live.

Seagrasses are important nurseries for marine life and help protect our coasts from storms and sea-level rise.

As important as the bacteria are, little was known about their makeup until now. Scientists at Clemson University have recently published a paper indicating that the bacteria are more taxonomically diverse than previously thought.

Normally, when seagrass decomposes, sulfide is produced which is toxic to seagrasses and clams. However, the bacteria oxidize the sulfide compounds to harmless sulfate, using the energy produced to turn carbon dioxide into sugars for the [clam](#).

"This whole symbiotic process enables the clams and bacteria to live in environments that are not necessarily hospitable," said Clemson biological sciences associate professor Barbara Campbell, whose research group studies how organisms interact with each other and how the microbiome—the community of microorganisms living in and on multicellular organisms—is involved in that interaction. "The more diverse the symbionts are, the healthier the ecosystem should be. Diversity matters."

Campbell's findings have important implications for fisheries and lobster farming, which require a healthy coastal ecosystem, including seagrass beds, in order to thrive.

Campbell and her students collected a sampling of *Ctena orbiculate* clams from the shallow waters at Sandy Creek Landing on Sugar Loaf Key in the far western Florida Keys. They also surveyed the

types of seagrass in the area, conducted physical and chemical surveys of the environment, and counted other clams.

In subsequent months, her team performed [next-generation sequencing](#) and analyses in her campus lab and at Clemson's Genomic and Bioinformatics Facility to characterize the taxonomic, genetic and functional composition of the symbiont bacterial communities.

"We looked at the DNA and RNA, in terms of what genes are present, the diversity of the genes and genomes, as well as the diversity of what is being expressed by the [bacteria](#) and the host," Campbell said.

According to Campbell, her research team plans to apply additional experiments to explore the diversity of the genes of both the clams and the microbes, which should provide more fundamental knowledge about the interactions between the two organisms.

"We can introduce a symbiont with a specific function versus a symbiont with a different function and then put the clams back into the seagrass and see what happens," said Campbell. They will see if more [diversity](#) leads to better seagrass health.

More information: Shen Jean Lim et al, Extensive Thioautotrophic Gill Endosymbiont Diversity within a Single *Ctena orbiculata* (Bivalvia: Lucinidae) Population and Implications for Defining Host-Symbiont Specificity and Species Recognition, *mSystems* (2019). [DOI: 10.1128/mSystems.00280-19](https://doi.org/10.1128/mSystems.00280-19)

Provided by Clemson University

Citation: Research indicates abundance of microbe diversity key to healthy coastal ecosystem

(2019, November 5) retrieved 20 April 2024 from <https://phys.org/news/2019-11-abundance-microbe-diversity-key-healthy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.