

Study of African animals illuminates links between environment, diet and gut microbiome

November 4 2019



New research analyzing the diets and microbiomes of 33 large-herbivore species in Kenya yields surprising findings about the interplay between animal evolution, behavior and the gut microbiome. Tyler Kartzinel collects a fecal sample at the Ol Jogi Conservancy in Central Kenya. The researchers analyzed more than

1,000 fecal samples. Credit: Robert Pringle, associate professor of ecology and evolutionary biology at Princeton University and senior author on the study.

In recent years, the field of microbiome research has grown rapidly, providing newfound knowledge—and newfound questions—about the microbes that inhabit human and animal bodies. A new study adds to that foundation of knowledge by using DNA analysis to examine the relationship between diet, the environment and the microbiome.

"Environmental change may influence what [animals](#) are eating, and as a consequence, influence their microbiome and health in a variety of ways that can only be understood in natural settings," said study lead author Tyler Kartzinel, an assistant professor of ecology and [evolutionary biology](#) at Brown University and a former postdoctoral researcher at Princeton University.

He added that the study's innovative DNA-based methods might ultimately provide new avenues to study and understand human microbiomes as well.

The research, published in the *Proceedings of the National Academy of Sciences* on Monday, Nov. 4, involved collecting and analyzing more than 1,000 samples of fecal material from 33 herbivore species—which ranged from diminutive dwarf antelopes to gigantic giraffes and elephants—in an African savanna.

To build on the findings of earlier studies, the research team—a collaboration of scientists in the ecology and evolutionary biology departments at Brown and Princeton, and colleagues from the botany department of the National Museums of Kenya—sought to study a wide variety of species by analyzing samples gathered from their natural

habitat. Much of the fieldwork was done at the Mpala Research Centre in Kenya, which is managed by Princeton University in partnership with the Smithsonian Institution, the National Museums of Kenya and the Kenya Wildlife Service.

"A fecal sample provides an amazing window into the biology of a wild animal, from what it eats to what bacteria live in its gut to what kinds of parasites it has," said Robert Pringle, an associate professor of ecology and evolutionary biology at Princeton and senior author on the study.

"We're just starting to tap into the potential of what forensic DNA-based approaches to wildlife ecology can teach us about these things that have historically been very difficult if not impossible to investigate."

After analyzing the DNA in the samples to infer the animals' diets and microbiomes, the researchers reached three main conclusions. Consistent with their expectations, they found that closely related species had similar microbiomes, and, to a lesser extent, similar diets. Their second finding was that species (and individual animals within a species) who consumed dissimilar diets tended to have dissimilar microbiomes.



New research analyzing the diets and microbiomes of 33 large-herbivore species in Kenya yields surprising findings about the interplay between animal evolution, behavior and the gut microbiome. An elephant forages on vegetation at the Mpala Research Centre in Kenya. Identifying the species of plants eaten by large wild herbivores such as elephants has historically been a major challenge for ecologists. Credit: Robert Pringle, associate professor of ecology and evolutionary biology at Princeton University and senior author on the study.

Lastly, the study found that animals whose diets underwent significant seasonal changes also tended to experience major seasonal changes in their microbiomes. But the team was surprised to find that the microbiomes of domesticated species such as cattle, sheep, goats, donkeys and camels tended to change more with the seasons than did the microbiomes of wild animals—even compared to the most closely

related wild animals that had similar diets.

Kartzinel suggested three possible factors that might explain the microbiome differences between livestock and wild animals: domestication, guidance of livestock to optimal food and water sources by human herders, and the protection of livestock in corrals at night. Together with his colleagues, he plans to explore precisely why some species' microbiomes are more sensitive to seasonal change.

Kartzinel noted that the methods employed in the study allowed for a deeper level of analysis. In the past, researchers studied microbiomes in one of two ways: Some researchers conducted "between-species" research, examining the microbiomes of a few animals representing different species—comparing, say, the microbiome of a sheep with that of a cow. Others conducted "within-species research," comparing the microbiomes of many more animals from the same species—across, say, different seasons. However, because Kartzinel and his colleagues analyzed DNA to measure diet from individual samples collected from many different species in the same environment, they were able to conduct between-species and within-species research simultaneously.

Previous between-species research tended to find that closely [related species](#) had more similar microbiomes, and within-species research found that seasons affected animals' microbiomes. The present study adds much more nuance to these findings.

"The work we're publishing begins to bridge those findings, making the relationships seem less binary," Kartzinel said. "Seasonal change isn't just present or absent, for example; we found instead that there's a gradient between the microbiomes that respond a lot and those that respond a little."

A variety of additional questions arise from the research. For example, is

seasonal sensitivity in the microbiome a sign of health or a sign of trouble?

"You can imagine animals changing their diets and microbiomes because they're good at adjusting to changes in the environment," Kartzinel said. "But you can also imagine them doing it because they're stressed out and just trying to survive as the environment changes."



A reticulated giraffe selects a mouthful of a Hibiscus plant at the Mpala Research Centre. The researchers analyzed plant DNA in animal fecal samples to quantify the diet composition for 33 herbivore species. Credit: Robert Pringle

More broadly, Kartzinel and his colleagues also hope to determine which

factor—diet or microbiome—tends to be more sensitive to the animal's environment. "The same plant can provide succulent fruit for animals to eat in one season, and only offer chewy twigs in the next—if the animal eats it in both seasons, our methods wouldn't register a change in that animal's diet, but the animal's gut microbiome would," he said.

He'd also like to explore experiments to determine the importance of diet and microbiome turnover for the health of wild animals.

"Is the sensitivity of an herbivore's microbiome going to help it keep a healthy diet in a changing world?" he asked. "Or do other adjustments and conditions take precedent as the animal makes decisions about how to survive? Maybe it's a little bit of both. We're talking about several endangered [species](#), and the livestock people depend on, so it's important to consider the possibilities."

If the microbiome does significantly influence animal health and behavior, Kartzinel said, then it could "affect entire food webs, communities and ecosystems, because it would determine survives and who does not. It's amazing to think about."

He added that in collaboration with a broader set of colleagues, the team is beginning to tap into the possibility that this research could affect humans.

"The biomedical world is really interested in figuring out whether—and how—we can manage the human gut microbiome to improve health, stress and nutrition," Kartzinel said. "Together with a whole suite of research approaches, we think these genetic methods of connecting [diet](#) and [microbiome](#) could provide an additional layer of information—for wildlife ecologists and biomedical researchers, too."

More information: Tyler R. Kartzinel et al., "Covariation of diet and

gut microbiome in African megafauna," *PNAS* (2019).
www.pnas.org/cgi/doi/10.1073/pnas.1905666116

Provided by Brown University

Citation: Study of African animals illuminates links between environment, diet and gut microbiome (2019, November 4) retrieved 24 April 2024 from
<https://phys.org/news/2019-11-african-animals-illuminates-links-environment.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.