

Amazon fires are causing glaciers in the Andes to melt even faster

November 29 2019, by Matthew Harris



Credit: AI-generated image ([disclaimer](#))

If you have turned on a TV or read the news during the past few months, you have probably heard of the widespread fires that wrought havoc on the Amazon rainforest this year. Fires occur in the rainforest every year, but the past 11 months saw the number of fires [increase by more than 70%](#) when compared with 2018, indicating a major acceleration in land

clearing by the country's logging and farming industries.

The smoke from the fires rose high into the atmosphere and could be seen from space. Some regions of Brazil became covered in thick smoke that closed airports and [darkened city skies](#).

As the rainforest burns, it releases enormous amounts of carbon dioxide, [carbon monoxide](#), and larger particles of so-called "[black carbon](#)" ([smoke and soot](#)). The phrase "enormous amounts" hardly does the numbers justice—in any given year, the burning of forests and grasslands in South America emits a whopping [800,000 tonnes of black carbon](#) into the atmosphere.

This truly astounding amount is almost double the black carbon produced by all combined energy use in Europe over 12 months. Not only does this absurd amount of smoke cause [health issues](#) and contribute to [global warming](#) but, as a growing number of scientific studies are showing, it also more directly contributes to the melting of [glaciers](#).

In a new paper published in the journal [Scientific Reports](#), a team of researchers has outlined how smoke from fires in the Amazon in 2010 made glaciers in the Andes melt more quickly.

When fires in the Amazon emit black carbon during the peak burning season (August to October), winds carry these clouds of smoke to Andean glaciers, which can sit higher than 5,000 metres above sea level.

Despite being invisible to the naked eye, black carbon particles affect the ability of the snow to reflect incoming sunlight, a phenomenon known as "albedo". Similar to how a dark-coloured car will heat up more quickly in direct sunlight when compared with a light-coloured one, glaciers covered by black carbon particles will absorb more heat, and

thus melt faster.

By using a computer simulation of how particles move through the atmosphere, known as [HYSPLIT](#), the team was able to show that smoke plumes from the Amazon are carried by winds to the Andes, where they fall as an invisible mist across glaciers. Altogether, they found that fires in the Amazon in 2010 caused a 4.5% increase in [water runoff](#) from Zongo Glacier in Bolivia.

Crucially, the authors also found that the effect of black carbon depends on the amount of dust covering a glacier—if the amount of dust is higher, then the glacier will already be absorbing most of the heat that might have been absorbed by the black carbon. Land clearing is one of the reasons that dust levels over South America [doubled](#) during the 20th century.

Glaciers are some of the most important natural resources on the planet. Himalayan glaciers provide drinking water for 240m people, and [1.9 billion rely on them for food](#). In South America, glaciers are crucial for water supply—in some towns, including Huaraz in Peru, [more than 85% of drinking water comes from glaciers during times of drought](#). However, these truly vital sources of water are increasingly under threat as the planet feels the effects of global warming. Glaciers in the Andes have been [receding](#) rapidly for the last 50 years.

The tropical belt of South America is predicted to become [more dry and arid](#) as the climate changes. A drier climate means more dust, and [more fires](#). It also means more droughts, which make towns more reliant on glaciers for water.

Unfortunately, as the above study shows, the fires assisted by dry conditions help to make these vital sources of water vanish more quickly. The role of black carbon in glacier melting is an exceedingly

complex process—currently, the climate models used to [predict the future melting of glaciers in the Andes](#) do not incorporate black carbon. As the authors of this new study show, this is likely causing the rate of glacial melt to be underestimated in many current assessments.

With communities reliant on glaciers for [water](#), and these same glaciers likely to melt faster as the climate warms, work examining complex forces like black [carbon](#) and albedo changes is needed more now than ever before.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Amazon fires are causing glaciers in the Andes to melt even faster (2019, November 29) retrieved 2 May 2024 from <https://phys.org/news/2019-11-amazon-glaciers-andes-faster.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.