

Antarctica likely to drive rapid sea-level rise under climate change

November 6 2019



Credit: CC0 Public Domain

Scientists from The Australian National University (ANU) have shown that ice melt from Antarctica drives rapid and high sea-level rise, offering a forewarning of what to expect under human-driven climate

change.

The researchers examined historical and new data from the 'last interglacial', which took place 125,000 to 118,000 years ago and saw [sea levels rise](#) up to 10 metres above current levels.

Interglacials are periods of warmer global temperatures that can last thousands of years.

The study, published in *Nature Communications*, shows that sea levels rose up to three metres per century, largely driven by [ice loss](#) in the Antarctic ice sheet.

Lead author, Professor Eelco Rohling, said that the last interglacial sea-rise was due to natural climate instabilities.

"These were smaller and slower than the human-caused [climate](#) disturbance of today," he said.

"Our study shows clearly that Antarctica, long thought a sleeping giant when it comes to sea-level rises, is in fact the key player.

"And it appears that it can change by large amounts on timescales that are highly relevant to society and in ways that would have profound effects on human infrastructure."

The study shows for the first time by how much ice loss in the last interglacial first took place in Antarctica, followed by Greenland.

Early Antarctic ice loss was caused by Southern Ocean warming at the onset of the interglacial. Next, the meltwater from Antarctica caused changes in global ocean circulation that resulted in northern polar warming and associated Greenland ice loss.

Co-lead author, Dr. Fiona Hibbert, said that in today's greenhouse-gas-driven [climate change](#), rapid atmospheric and oceanic warming happens in both polar regions at the same time.

"This drives simultaneous ice-loss in Antarctica and Greenland," Dr. Hibbert said.

"But, what's vital to remember is that today's climatic disturbance is greater and develops faster than that of the last interglacial.

"As a result, rates of sea-level rise may develop over the next several centuries that are even higher than those found for the interglacial we have studied."

The study was jointly led Professor Rohling and Dr. Hibbert from ANU, with from colleagues from Australia, Norway, Spain, the United States, and Germany.

More information: Eelco J. Rohling et al. Asynchronous Antarctic and Greenland ice-volume contributions to the last interglacial sea-level highstand, *Nature Communications* (2019). [DOI: 10.1038/s41467-019-12874-3](#)

Provided by Australian National University

Citation: Antarctica likely to drive rapid sea-level rise under climate change (2019, November 6) retrieved 18 April 2024 from <https://phys.org/news/2019-11-antarctica-rapid-sea-level-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.