


ATLAS Experiment probes the quark-gluon plasma in a new study of photo-produced muon pairs

November 27 2019



The distributions of k_{\perp} in different centrality intervals, where the percentage values indicate the fraction of events in a given centrality class starting from the most central events (0-5%). The bottom left panel corresponds to the UPC

events. The p_T of both muons in the pair is required to be larger than 4 GeV . The distributions are normalized such that the integral of the distributions in each centrality class is equal to the ratio of dimuon pairs in that centrality class to the total number of dimuon pairs across all centralities (including the UPC events). The red lines indicate parameterized fits to the distributions. Credit: ATLAS Collaboration/CERN

At the [Large Hadron Collider](#) (LHC) at CERN, the electromagnetic fields of Lorentz-contracted lead nuclei in heavy-ion collisions act as intense sources of high-energy photons, or particles of light. This environment allows particle physicists to study photon-induced scattering processes, [which can not be studied elsewhere](#).

A key process examined by [ATLAS Experiment](#) physicists involves the annihilation of photons into pairs of oppositely charged muons. Such electromagnetic "two-photon processes" are typically studied in "ultra-peripheral collisions" (UPC), where the transverse separation between the colliding [lead nuclei](#) is greater than the sum of their radii, resulting in no direct strong interactions between the colliding nuclei. This provides a [clean environment](#) for the study of electromagnetic interactions at high energy and intensity. However, these two-photon processes are also present in collisions where the two nuclei overlap ("central collision events") and produce quark-gluon plasma. The produced muons can, in principle, interact with the charges in the plasma, making the muon pairs created in two-photon processes a potentially valuable probe of the electromagnetic fields in the plasma.

The ATLAS Collaboration recently released a [new, comprehensive measurement](#) of the distributions of muon pairs from two-photon annihilation processes, in UPC and non-UPC collision events. The measurement utilizes the large dataset recorded during the 2015 and

2018 heavy-ion runs of the LHC.

ATLAS physicists found that the distributions of muon pairs varied systematically depending on the "centrality" of the collision (a measure of how head-on two nuclei collide). This behavior is quantified by the observable k_{\perp} which represents the transverse momentum of the dimuon pair perpendicular to the [muon](#) directions. The figure shows the [distribution](#) of several different centrality classes, ranging from UPC events to central collision events.

A significant change in the distributions is observed from UPC to peripheral to central collision events. In particular, for the UPC events, the two muons are most likely to be produced back-to-back, leading to the k_{\perp} distributions peaking at $k_{\perp} = 0$ MeV. However, in more central collisions with hadronic interactions, the two muons are more likely to have a slight shift from being purely back-to-back, resulting the k_{\perp} distributions to have a most probable value larger than zero. The most probable value of k_{\perp} shifts, depending on the centrality of the [collision](#) event, from $k_{\perp} = 0$ MeV in UPC events to $k_{\perp} = 36 \pm 1$ MeV in the 0-5% most central collisions.

These measurements provide new insight into the possible interaction of the outgoing muons with electromagnetic charges or fields present in the quark-gluon plasma. However, recent calculations suggest that effects similar to those seen in the data may result from a combination of the initial state broadening of the photon transverse momenta and from the production process itself. Future analyses and additional measurements are needed to establish the mechanism(s) responsible for the features observed in the data.

More information: Measurement of non-exclusive dimuon pairs produced via $\gamma\gamma$ scattering in lead-lead collisions at 5.02 TeV with the ATLAS detector (ATLAS-CONF-2019-051):

[atlas.web.cern.ch/Atlas/GROUPS ... ATLAS-CONF-2019-051/](https://atlas.web.cern.ch/Atlas/GROUPS/CONFERENCE/notes/ATLAS-CONF-2019-051/)

Provided by ATLAS Experiment

Citation: ATLAS Experiment probes the quark-gluon plasma in a new study of photo-produced muon pairs (2019, November 27) retrieved 30 April 2024 from

<https://phys.org/news/2019-11-atlas-probes-quark-gluon-plasma-photo-produced.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--