

Carbon dots make calcium easier to track



November 12 2019, by Zhang Nannan

Illustration of Ca2+ detection by EGTA-modified CDs. Credit: SIBET

In hospitals, doctors often advise patients to take calcium supplements. But does the calcium get into the cells that need it? Until recently, it's been hard to tell.

But now, Prof. Dong Wenfei's research group from the Suzhou Institute of Biomedical Engineering and Technology (SIBET) has developed a new type of fluorescent carbon dot that can effectively detect <u>calcium</u> levels in cells.

Calcium is essential to life. Calcium ions regulate a variety of important cellular functions such as differentiation, proliferation, growth and gene transcription.

In addition, studies have reported that calcium status in cancer cells is



associated with tumorigenesis, metastasis and angiogenesis. Other studies have also shown that abnormal changes in intracellular Ca²⁺ concentration are associated with the occurrence and development of Parkinson's disease, Huntington's disease and Alzheimer's disease. Thus, detection of intracellular calcium levels is very important.

Conventional methods for calcium ion detection such as atomic absorption spectroscopy (AAS) and <u>nuclear magnetic resonance</u> (NMR) require expensive equipment and tedious sample preparation, limiting their applications in intracellular detection. Therefore, developing affordable, efficient, sensitive and simple techniques for detecting Ca²⁺ ions in cells is a priority. Fortunately, fluorescence detection meets these requirements.

As a new fluorescent probe, carbon dots (CDs) have been widely used in ion and small-molecule detection due to their extraordinary performance, which includes water solubility, excellent stability, nontoxicity, adjustable fluorescence and high quantum yield.

In this study, the researchers used an innovative approach to preparing functional CDs. They first synthesized the CDs from citric acid and ethylenediamine using a hydrothermal method. Then they modified the surface of the CDs with EGTA by reheating the solution.

This is a simple and convenient chemical preparation method, compared with the traditional cross-linking procedure, because all reactions are carried out in the same autoclave. The EGTA-modified CDs are the final product; no secondary purification and separation is required.

This work entitled "Two-Step Hydrothermal Preparation of Carbon Dots for Calcium Ion Detection" has been published in *ACS Applied Materials & Interface*.



The CDs exhibit bright blue fluorescence. As calcium concentration increases, the fluorescence intensity drops sharply.

Cellular experiments show that CDs are nontoxic and have good biocompatibility, suggesting they may be a useful probe for intracellular Ca^{2+} detection.

In the future, researchers expect to develop fluorescent carbon dots in the near-infrared band with strong penetration. The distribution of <u>calcium ions</u> in internal organs could then be detected nondestructively.

More information: Juan Yue et al, Two-Step Hydrothermal Preparation of Carbon Dots for Calcium Ion Detection, *ACS Applied Materials & Interfaces* (2019). DOI: 10.1021/acsami.9b13737

Provided by Chinese Academy of Sciences

Citation: Carbon dots make calcium easier to track (2019, November 12) retrieved 2 May 2024 from <u>https://phys.org/news/2019-11-carbon-dots-calcium-easier-track.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.