

New Cretaceous mammal fossil sheds light on evolution of middle ear

November 27 2019



Reconstruction of *Jeholbaatar kielanae*. Credit: XU Yong

Researchers from the Institute of Vertebrate Paleontology and Paleoanthropology (IVPP) of the Chinese Academy of Sciences and the American Museum of Natural History (AMNH) have reported a new species of multituberculate—a type of extinct Mesozoic rodent—with well-preserved middle ear bones from the Cretaceous Jehol Biota of China. The findings were published in *Nature* on November 27.

The new [mammal](#), *Jeholbaatar kielanae*, has a middle ear that is distinct from those of its relatives. Wang Yuanqing and Wang Haibing from IVPP, along with Meng Jin from AMNH, proposed that the evolution of its auditory apparatus might have been driven by specialization for feeding.

Fossil evidence shows that postdentary bones were either embedded in the postdentary trough on the medial side of the dentary or connected to the dentary via an ossified Meckel's cartilage in early mammals, prior to their migration into the cranium as seen in extant mammals.

Detachment of the mammalian [middle ear bones](#) from the dentary occurred independently at least three times. But how and why this process took place in different clades of mammals remains unclear.

The *Jeholbaatar kielanae* specimen was discovered in the Jiufotang Formation in China's Liaoning Province (Jehol Biota). It displays the first well-preserved middle-ear bones in multituberculates, providing solid evidence of the morphology and articulation of these bony elements, which are fully detached from the dentary.

It reveals a unique configuration with more complete components than those previously reported in multituberculates. The new fossil reveals a transitional stage in the evolution of the surangular—a "reptilian" jawbone.

In light of current evidence, scientists argue that the primary (malleus-incus) and secondary (squamosal-dentary) jaw joints co-evolved in allotherians, allowing a distinct palinal (anteroposterior) jaw movement while chewing.

Detachment of the auditory apparatus of the middle ear would have gained higher [selective pressure](#) in order to increase feeding efficiency, suggesting that evolution of the middle ear was probably triggered by functional constraints on the feeding apparatus in allotherians.

More information: Cretaceous fossil reveals a new pattern in mammalian middle ear evolution, *Nature* (2019). [DOI: 10.1038/s41586-019-1792-0](#) , [nature.com/articles/s41586-019-1792-0](https://www.nature.com/articles/s41586-019-1792-0)

Provided by Chinese Academy of Sciences

Citation: New Cretaceous mammal fossil sheds light on evolution of middle ear (2019, November 27) retrieved 2 May 2024 from <https://phys.org/news/2019-11-cretaceous-mammal-fossil-evolution-middle.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--