

## Study shows cover crops and perennials do not necessarily increase carbon storage in soil

November 11 2019, by Fred Love



Corn and reconstructed prairie plots at a long-term ISU field experiment. Credit: Steven Hall



The ability of cover crops to stimulate microbes deep in the soil of farm fields leads to significant gains in water quality but does not necessarily increase the capacity of soil to store carbon, according to a recently published study from Iowa State University scientists.

The study, published in the academic journal *Global Change Biology Bioenergy*, analyzed <u>soil samples</u> that had undergone a wide range of long-term treatments. Steven Hall, an assistant professor of ecology, evolution and organismal biology and corresponding author of the study, said the research delves deep below the surface to explore how <u>cover</u> <u>crops</u> and perennial prairie plants affect soil microbial activity.

The research found winter rye, a plant commonly used as a cover crop, provides a source of carbon that stimulates microbes as much as a meter under the surface of the soil. That stimulation increases the microbes' demand for nutrients. The process of consuming soil nutrients keeps those nutrients from running into waterways, which is an important environmental benefit.

However, the microbes may digest much of the carbon provided by the cover crop, converting it into <u>carbon dioxide</u>, which is released to the atmosphere. This means that the cover-crop carbon is not necessarily stored, or sequestered, where it could help offset climate change.

The findings may surprise some who assume having more plants in the soil means increased carbon storage, Hall said.

"Cover crops and perennials provide key benefits for <u>water quality</u>, but I wouldn't hang my hat on rapid carbon sequestration benefits," he said. "We've found a tradeoff. Greater plant growth doesn't necessarily mean gains in carbon sequestration if microbial activity also increases."

The study analyzed 96 soil <u>core samples</u> taken from a 10-year field



study. Some of the fields underwent continuous corn planting, some underwent corn-soybean rotation, some had cover crops following continuous corn and some had reconstructed prairies. The core samples allowed the researchers to figure out how deeply cover crops and perennial plants intensify microbial activity, and to see what carbon sources microbes were digesting.

"Despite their notable environmental benefits, neither unfertilized perennials nor cover crops necessarily promote rapid <u>soil carbon</u> sequestration relative to conventional annual bioenergy systems because of concomitant increases in decomposition," the study reported.

**More information:** Chenglong Ye et al. Mechanisms underlying limited soil carbon gains in perennial and cover-cropped bioenergy systems revealed by stable isotopes, *GCB Bioenergy* (2019). DOI: 10.1111/gcbb.12657

Provided by Iowa State University

Citation: Study shows cover crops and perennials do not necessarily increase carbon storage in soil (2019, November 11) retrieved 27 June 2024 from <u>https://phys.org/news/2019-11-crops-perennials-necessarily-carbon-storage.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.