


# Hot electrons harvested without tricks

November 15 2019

---



A set up for ultrafast spectroscopy, as used in the study. Credit: Maxim Pchenitchnikov, University of Groningen

Semiconductors convert energy from photons (light) into an electron current. However, some photons carry too much energy for the material to absorb. These photons produce "hot electrons," and the excess energy of these electrons is converted into heat. Materials scientists have been

looking for ways to harvest this excess energy. Scientists from the University of Groningen and Nanyang Technological University (Singapore) have now shown that this may be easier than expected by combining a perovskite with an acceptor material for hot electrons. Their proof of principle was published in *Science Advances* on 15 November.

In photovoltaic cells, semiconductors will absorb [photon energy](#), but only from photons that have the right amount of energy: too little, and the photons pass right through the material; too much, and the [excess energy](#) is lost as heat. The right amount is determined by the bandgap: the difference in [energy levels](#) between the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO).

## Nanoparticles

"The excess energy of hot electrons produced by the high-energy photons is very rapidly absorbed by the material as heat," explains Maxim Pshenichnikov, professor of ultrafast spectroscopy at the University of Groningen. To fully capture the energy of hot electrons, materials with a larger bandgap must be used. However, this means that the hot electrons should be transported to this material before losing their energy. The current general approach to harvesting these electrons is to slow down the loss of energy, for example, by using nanoparticles instead of bulk material. "In these nanoparticles, there are fewer options for the electrons to release the excess energy as heat," explains Pshenichnikov.

Together with colleagues from Nanyang Technological University, where he was a visiting professor for the past three years, Pshenichnikov studied a system in which an organic-inorganic hybrid perovskite semiconductor was combined with the organic compound bathophenanthroline (bphen), a material with a large bandgap. The

scientists used laser light to excite electrons in the perovskite and studied the behavior of the hot electrons that were generated.

## Barrier

"We used a method called pump-push probing to excite electrons in two steps and study them at femtosecond timescales," explains Pshenichnikov. This allowed the scientists to produce electrons in the perovskites with energy levels just above the bandgap of bphen, without exciting electrons in the bphen. Therefore, any hot electrons in this material would have come from the perovskite.

The results showed that hot electrons from the perovskite semiconductor were readily absorbed by the bphen. "This happened without the need to slow down these electrons, and moreover, in bulk material. So without any tricks, the hot electrons were harvested." However, the scientists noticed that the energy required was slightly higher than the bphen bandgap. "This was unexpected. Apparently, some extra energy is needed to overcome a barrier at the interface between the two materials."

Nevertheless, the study provides a proof of principle for the harvesting of [hot electrons](#) in bulk [perovskite](#) semiconductor material.

Pshenichnikov says, "The experiments were performed with a realistic amount of [energy](#), comparable to visible light. The next challenge is to construct a real device using this combination of materials."

**More information:** "Hot carrier extraction in CH<sub>3</sub>NH<sub>3</sub>PbI<sub>3</sub> unveiled by pump-push-probe spectroscopy" *Science Advances* (2019).

[dx.doi.org/10.1126/sciadv.aax3620](https://doi.org/10.1126/sciadv.aax3620) ,  
[advances.sciencemag.org/content/5/11/eaax3620](https://advances.sciencemag.org/content/5/11/eaax3620)

Provided by University of Groningen

Citation: Hot electrons harvested without tricks (2019, November 15) retrieved 20 March 2024 from <https://phys.org/news/2019-11-hot-electrons-harvested.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.