


Massive photons in an artificial magnetic field

November 12 2019




The dependence of the energy (vertical axis) on the angle for polarized light reflected from birefringent optical cavity (horizontal axis). Credit: M. Krol, UW Physics

An international research collaboration from Poland, the UK and Russia has created a two-dimensional system—a thin optical cavity filled with liquid crystal—in which they trapped photons. As the properties of the cavity were modified by an external voltage, the photons behaved like massive quasiparticles endowed with a magnetic moment, called "spin," under the influence of an artificial magnetic field. The research has been published in *Science* on Friday, 8 November 2019.

The world around us has one temporal and three spatial dimensions. Physicists studying condensed matter have long been dealing with systems of lower dimensionality—two-dimensional (2-D) quantum wells, one-dimensional (1-D) quantum wires and zero-dimensional (0-D) quantum dots. 2-D systems have found the widest technical applications—it is thanks to the reduced dimensions that efficient LEDs and laser diodes, fast transistors in integrated circuits, and WiFi radio amplifiers operate. Trapped electrons in two dimensions can behave completely differently than [free electrons](#). For example, in graphene, a two-dimensional carbon structure with honeycomb symmetry, electrons behave like massless objects, i.e. [light particles](#) called photons.

Electrons in a crystal interact with each other and with the [crystal lattice](#), creating a complex system whose description is possible thanks to the introduction of the concept of so-called quasiparticles. Properties of these quasiparticles, including electric charge, [magnetic moment](#) and

mass, depend on the symmetry of the crystal and its spatial dimension. Physicists can create materials with reduced dimensions, discovering "quasi-universes" full of exotic quasiparticles. The massless electron in two-dimensional graphene is such an example.




Tomography of circularly polarized light reflected from an optical cavity filled with liquid crystal. Credit: M. Krol, UW Physics

These discoveries inspired researchers from the University of Warsaw, the Polish Military University of Technology, the Institute of Physics of the Polish Academy of Sciences, the University of Southampton and the Skolkovo Institute near Moscow, to study light trapped in two-dimensional structures—optical cavities.

The authors of the *Science* paper created an optical cavity in which they trapped photons between two mirrors. The original idea was to fill the cavity with a liquid crystal material that acts as an optical medium. Under the influence of an external voltage, molecules of this medium can rotate and change the optical path length. Because of this, it was possible to create standing waves of light in the cavity, whose energy (frequency of vibrations) was different when the electric field of the wave (polarization) was directed across the molecules and different for polarization along their axis (this phenomenon is called optical anisotropy).

During the research, conducted at the University of Warsaw, the unique behavior of photons trapped in the cavity was found as they behaved like mass-bearing quasiparticles. Such quasiparticles have been observed before, but they were difficult to manipulate because the light does not react to electric or magnetic fields. This time, it was noted that as the optical anisotropy of the liquid crystal material in the cavity was changed, the trapped photons behaved like quasiparticles endowed with a magnetic moment, or a "spin" in an "artificial magnetic field." Polarization of the electromagnetic wave played the role of "spin" for light in the cavity. The behavior of light in this system is easiest to explain using the analogy of the behavior of electrons in condensed matter.



The scheme of the experiment - circular polarization of light (marked in red and blue) transmitted through a cavity filled with liquid crystal depending on the

direction of propagation. Credit: M. Krol, UW Physics

The equations describing the motion of photons trapped in the [cavity](#) resemble the equations of motion of electrons with spin. Therefore, it was possible to build a photonic system that perfectly imitates electronic properties and leads to many surprising physical effects such as topological states of light.

The discovery of new phenomena related to the entrapment of [light](#) in optically anisotropic cavities may enable the implementation of new optoelectronic devices, e.g. optical neural networks and perform neuromorphic calculations. There is particular promise to the prospect of creating a unique quantum state of matter—the Bose Einstein condensate. Such a condensate can be used for quantum calculations and simulations, solving problems that are too difficult for modern computers. The studied phenomena will open up new possibilities for technical solutions and further scientific discoveries.

More information: Katarzyna Rechcińska et al, Engineering spin-orbit synthetic Hamiltonians in liquid-crystal optical cavities, *Science* (2019). [DOI: 10.1126/science.aay4182](https://doi.org/10.1126/science.aay4182)

Provided by University of Warsaw

Citation: Massive photons in an artificial magnetic field (2019, November 12) retrieved 2 May 2024 from <https://phys.org/news/2019-11-massive-photons-artificial-magnetic-field.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.