

Mechanisms of soft tissue and protein preservation in Tyrannosaurus rex

November 12 2019, by Thamarasee Jeewandara



A fossil specimen of T. rex (AMNH 5027) on display at the American Museum of Natural History, in 2019. Credit: National Museum of Natural History, New York City. amnh.org/exhibitions/permanent/saurischian-dinosaurs/tyrannosaurus-rex.

The existing notion that soft tissue architectures and native proteins can be preserved across geological time is controversial since methods of such preservation remain to be investigated and well-defined. In a new



study, Elizabeth M. Boatman and colleagues at the departments of Engineering, Paleontology, Biological Science, Materials and Engineering and the Advanced Light Source in the U.S., tested crosslinking mechanisms for preserved tissue architecture. They used two non-enzymatic, structural protein mechanisms, Fenton chemistry and glycation to demonstrate their possible contributions to preserve blood vessel structures recovered from the cortical bone of *Tyrannosaurus rex* (*T. rex*; USNM 555 000, formerly MOR 555). They demonstrated the endogeneity (randomness) of the fossil vessel tissues and the presence of type I collagen in the outermost vessel layers using imaging, diffraction, spectroscopy and <u>immunohistochemistry</u>.

They derived data from synchrotron Fourier transform infrared (SR-FTIR) studies on the *T. rex* vessels to analyze their crosslinking character and compared them with control chicken samples treated similarly with the two techniques. The researchers provided X-ray microprobe analyses of the chemical state of the fossil tissues to support <u>vessel</u> preservation of *T. rex*, as observed using the methods of investigation. Boatman et al. propose that the observed tissue stabilizing crosslinks will play an important role to preserve additional microvascular tissues in skeletal elements from the <u>Mesozoic era</u>. The work is now published on *Scientific Reports*.

Paleontologists have recovered hollow, pliable and transparent vessellike structures from skeletal elements of fossil vertebrates including nonavian dinosaurs and applied many techniques to identify their endogenous proteins such as <u>collagen</u> and <u>elastin</u>. Researchers had used mass-spectroscopy sequencing to identify isolated vessels recovered from non-avian dinosaurs to support the presence of vertebrate-specific vascular proteins in the past. For example, they documented the hallmark 67-nanometer-banding pattern typical for type I collagen after liberating the protein via demineralization, followed by additional studies to verify the presence of type I collagen in vascular canals of a



sauropod dinosaur rib from approximately 190 million years ago using <u>FTIR and Raman analysis</u>. While research teams had developed a variety of methods to explain unexpected preservation, experimental testing of proposed mechanisms remain to be conducted routinely and broadly.

In the present work, Boatman et al. identified and tested the possible contribution of a set of experiments to preserve the vessel-like architecture of the compact bone of a *Tyrannosaurus rex* fossil. They expect the work to lay a possible foundation for additional studies on preserving soft tissues recovered from the Mesozoic or more recent fossils. The walls of vertebrate blood vessel contain three distinct layers including the tunica intima (innermost), tunica media and tunica externa (outermost layer). Due to their unique molecular compositions, scientists can differentiate the constituents morphologically and chemically. For example, elastin is a helical protein specific to vertebrates that offers resistance to pressure changes in the vascular walls. Collagen is also vertebrate-specific and constitutes a predominant fraction of blood vessels to serve as their structural foundation. Since elastin and collagen contain hallmark features identifiable at the molecular structure and composition, Boatman et al. proposed to study the two proteins within the remnant dinosaur vessels.





LEFT: SR-FTIR analysis. Amide I sub-band localization of untreated and treated chicken type I collagen in SR-FTIR spectra. Sub-bands (β -sheet, ~1633 cm-1; triple-helix, ~1658–1660 cm-1; intermolecular, ~1683–1690 cm-1) are indicated in the figures. Red traces denote second derivatives of experimental curves. Although the intermolecular sub-band typically presents at lower wavenumber, the identified value was the nearest local minimum in each of the second derivative traces and consistently appears across all samples; therefore, in this sample, the intermolecular sub-band was indexed at 1697–1699 cm-1. RIGHT: SEM images of USNM 555000 cortical bone. (a) Fracture surface showing clear features of osteons (o) predominantly in longitudinal section, osteocyte lacunae (ol; in dashed white circles), and fine texture consistent with mineralized collagen fibers in bone. Back-scattered (BSE) image. (b), Polished (1200 grit) transverse section (BSE image) showing clear features of osteons and osteocyte lacunae. Mineral infilled osteons (white arrows) yield highly altered vessel structures, which were readily eliminated from SAXS, FTIR, and TEM analysis by careful preparation (sedimentation, washing, selection under microscope). Cracks are due to humidity/pressure changes and are an artefact of preparation. (c), Polished (1200 grit) transverse section (secondary electron [SE] image) showing clear features of osteons and osteocyte lacunae. (d), Highly magnified SE image of an osteon, showing fibrous texture at edges (white arrow), which was commonly observed in non-mineral infilled osteons in this specimen. This thin, fibrous coating inside the osteon structure is proposed to be the hollow, pliable vessel structures. Credit: Scientific Reports, doi: 10.1038/s41598-019-51680-1

The research team hypothesized the contribution of early <u>diagenetic</u> (physical and chemical) processes to the survival of *T. rex* microvasculature from deep-time. To test this, Boatman et al. first conducted SR-FTIR analysis to understand crosslink character in their control sample of chicken type I collagen protein. They induced crosslinks in the protein using <u>Fenton reagent or ion-catalyzed glycation</u> <u>techniques</u> followed by the use of transmission SR-FTIR to test each tissue. They observed the intramolecular crosslinks formed in the



chicken tissues to be immature due to their lack of exposure to pathways necessary to form <u>intermolecular crosslinks or advanced glycation</u> <u>endproducts (AGEs)</u>.

To test the *T. rex* vessel architecture for endogenous proteins, the scientists liberated three types of vessels from a demineralized *T. rex* cortical bone. They then used visible light microscopy (VLM) to characterize them as:

- 1. Extensive, brown-hued pliable networks
- 2. Fragmented opaque structures
- 3. Fragmented semi-translucent structures

They coupled <u>energy-dispersive X-ray spectroscopy</u> (EDS) with <u>scanning electron microscopy</u> (SEM) as well as <u>micro-focused X-ray</u> <u>fluorescence</u> (μ XRF) spectroscopy to confirm the differences observed in the tissue samples of varying composition. The team focused on the pliable vessel networks due to their similarly to existing bone tissue, which presumably maintained minimal alteration.







LEFT: Microscopy images of T. rex vascular tissue and associated analysis of fibrillar collagen banding. (a) Transmitted VLM of T. rex soft tissue shows an extensive network of hollow, pliable, vascular structure and typical brown hue. (b) SEM image of the surface of a vessel. (c) Magnified image of (b) detailing features consistent with collagen fibre bundles (collagen fibril, "f"; collagen fibre, "CF"). Average fibril width was measured as 110 nm, and average fibre width, 1.0 µm. (d) TEM image of fibrous features observed in a longitudinal vessel cross-section. Intensity profiles of banded texture in (e) boxes 1 and 2 in c and (f) boxes 3, 4, 5 in (d) with example peak-to-peak distances (SEM average, ~74 nm; TEM, ~56 nm) called out in red. RIGHT: Images and chemical analysis of the organic, brown-hued Tyrannosaurus rex vessels and mineralized casts of the vessel system liberated upon tissue demineralization. (a), Optical microscopy image of typical vessel fragments liberated from T. rex. (b), Magnified image of white rectangle in a, depicting brown-hued, pliable, organic vessel (ov) fragments and two types of mineralized casts: opaque forms and semitranslucent forms. (c), SEM image of similar vessel fragments. (d), Magnified image of white rectangle in c, with three vessel fragment types identified and locations of EDS analysis in e explicitly identified. (e), EDS analysis identified the opaque vessel casts as an iron oxide form (bottom, red) and the semitranslucent vessel casts as BaSO4 (top, blue). Credit: Scientific Reports, doi: 10.1038/s41598-019-51680-1

When Boatman et al. studied the pliable *T. rex* vessels using SEM, they observed fibrous structures across their outermost surface. The combined features were consistent with those observed in extant vessels liberated from cortical bone and <u>with fibrillar collagen</u>. The team analyzed the SR-FTIR spectrum of *T. rex* vessels to detect the dominant bands observed in both treated extant and ancient tissues. Notably, the amide I band for the dinosaur tissue was located at a predominant α -helix structure consistent with mature (crosslinked) fibrillar collagen. The research team then conducted <u>immunohistochemistry</u> (IHC) studies



to identify protein-specific epitopes of the structural proteins elastin and type I collagen.

The scientists raised antibodies against all components of the extant vasculature to observe positive binding in the dinosaur vessel walls. Using a fluorescent filter, they captured the localization and distribution of antibody-antigen complexes (green fluorescence). The response of the dinosaur vessels to actin antibodies appeared as a thin and evenly distributed layer. Antibodies raised against the muscle protein tropomyosin appeared with greater intensity on the vessel walls. The dinosaur vessels also indicated the presence of type I collagen antibodies, although elastin antibodies showed greater intensity. The two proteins were good targets for fossil studies due to high evolutionary conservation in certain regions. They did not observe reactivity of dinosaur vessels to antibodies against bacterial peptidoglycan (indicating no microbial contamination).





T. rex tissues exhibit positive antibody binding to protein components of extant vascular tissue. (a,c,e,g,i,k,m,o) Are composite images in which fluorescence corresponding to antibody-antigen complexes is overlain upon VLM images of vessel sections, with adjacent images (b,d,f,h,j,l,n,p) captured using a fluorescent filter. (a–d) No spurious binding was observed for negative controls in which vessels were exposed to secondary antibodies raised against the host species of all other antibodies used, i.e., mouse (a,b) and rabbit (c,d). (e,f) Positive binding of dinosaur vessels to actin antibodies can be seen in thin, evenly distributed layers, and (g,h) more broadly distributed binding is apparent for muscle tropomyosin antibodies. Antibodies to both (i,j) type I collagen and (k,l) elastin bind positively to these T. rex vessels. (m,n) Antibodies raised against ostrich haemoglobin exhibit comparatively lower binding intensity. (o,p) No reactivity of dinosaur vessels to antibodies against bacterial peptidoglycan was observed (indicating no contamination). Credit: *Scientific Reports*, doi: 10.1038/s41598-019-51680-1



Boatman et al. tested *T. rex* vessel structures to understand if postmortem structural protein crosslinking enhanced their resistance to degradation or diagenetic alterations. For this, they focused on fibrillar collagen using SR-FTIR transmission spectra to suggest post-mortem crosslinking during the process of <u>tissue architecture</u> preservation. These spectral features were previously recorded but not discussed with <u>early</u> <u>Jurassic sauropodomorphs</u> and <u>cretaceous bones</u>. The scientists then treated bulk *T. rex* tissue with <u>sodium borohydride</u> (NaBH₄) to reduce carbonyl groups within immature crosslinks and increase the non-peptide carbonyl absorption intensity. The carbohydrate absorption bands in the *T. rex* tissue were consistent with AGEs (advanced glycation endproducts). After treatment, the data suggested that *T. rex* tissues possessed both intramolecular and intermolecular crosslink types.

When the scientists mapped the elements in the tissue using μ XRF, they revealed iron (Fe) as the only metal concentrated within the dinosaur vessel tissues while recording barium (Ba) within the semi-translucent vessel casts. Using extended micro X-ray absorption near-edge structure microscopy, they observed Fe³⁺ embedded in the vessel walls. The researchers showed the presence of finely crystalline goethite (α -FeO(OH)); a mineral previously detected in vascular tissues recovered from two diverse dinosaur specimens.





TOP: SR-FTIR analysis of T. rex vascular tissue, NaBH4 reduced T. rex vascular tissue, chicken type I collagen without treatment, and chicken type I collagen treated with Fenton reagent and iron-catalyzed glycation. (a,b) Average FTIR spectra in the non-peptide carbonyl and protein amide I regions for all five samples. (a) Significant reduction in the non-peptide carbonyl band follows treatment of T. rex vascular tissue with NaBH4, which reduces (immature) peptide crosslinks. The blue-shifted Amide I band of the dinosaur tissue, Fenton reagent-treated chicken type I collagen, and Fe-catalyzed glycation-treated chicken type I collagen indicate increasing α -helix structure (~1660 cm-1) as the higher-energy triple-helix and intermolecular sub-bands increasingly predominate the spectra. The development of aldehydic carbonyl, ketoaldehyde, and/or immature ketoimine bands in both treated chicken tissues is consistent with the strong carbonyl band in the dinosaur tissue. BOTTOM: Micro-X-ray fluorescence (u-XRF) mapping of organic and mineralized Tyrannosaurus rex



vessel tissues reveal presence of multiple metals. S, Fe, and Ba were the primary elements identified in the vessel samples. Composite maps of two (a,b show Ba and Fe) and three (c,d show S, Ba, and Fe) atomic species show that Fe is predominantly associated with the organic vessel tissues, whereas S and Ba are mainly associated with the semi-translucent mineralized vessel casts. Credit: *Scientific Reports*, doi: 10.1038/s41598-019-51680-1

In this way, Elizabeth M. Boatman and colleagues demonstrated the presence of vertebrate-species endogenous proteins within soft tissue dinosaur structures. This included the presence of type I collagen consistent with the vasculature in extant vertebrates. The data supported a two-step mechanism that stabilized biomolecules and vessel architecture after the death of the organism, to promote their preservation within skeletal elements. The team hypothesized that iron-mediated Fenton and glycation pathways may have contributed to enhanced *T. rex* tissue longevity of elastin and fibrillar collagen within and around blood vessels. Both processes could be catalyzed by transition metal species such as iron to define the central role of Fe observed in structural protein crosslinking. The formation of iron oxyhydroxide precipitates in the work fully supported this idea.

The data represent the first comprehensive chemical and molecular characterization of vascular tissues recovered from *T. rex* specimen USNM 555000. The results shed light on the possible processes of fossilization at the molecular level. The researchers envision the demonstrated techniques will contribute to the development of comprehensive mechanisms to consistently retain vascular <u>tissue</u> survival from deep time.

More information: Elizabeth M. Boatman et al. Mechanisms of soft tissue and protein preservation in Tyrannosaurus rex, *Scientific Reports*



(2019). DOI: 10.1038/s41598-019-51680-1

Mary H. Schweitzer et al. A role for iron and oxygen chemistry in preserving soft tissues, cells and molecules from deep time, *Proceedings of the Royal Society B: Biological Sciences* (2013). DOI: 10.1098/rspb.2013.2741

Sergio Bertazzo et al. Fibres and cellular structures preserved in 75-million–year-old dinosaur specimens, *Nature Communications* (2015). DOI: 10.1038/ncomms8352

© 2019 Science X Network

Citation: Mechanisms of soft tissue and protein preservation in Tyrannosaurus rex (2019, November 12) retrieved 28 April 2024 from <u>https://phys.org/news/2019-11-mechanisms-soft-tissue-protein-tyrannosaurus.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.