

Microplastics found in oysters, clams on Oregon coast, study finds

November 12 2019



Credit: CC0 Public Domain

Tiny threads of plastics are showing up in Pacific oysters and razor



clams along the Oregon coast—and the yoga pants, fleece jackets, and sweat-wicking clothing that Pacific Northwesterners love to wear are a source of that pollution, according to a new Portland State University study.

Britta Baechler, a Ph.D. student in PSU's Earth, Environment and Society program, and Elise Granek, a professor of environmental science and management, looked at what variables predict <u>microplastic</u> concentrations in Pacific oysters and razor clams—organisms that have commercial, recreational and cultural importance in Oregon.

On average, the researchers found 11 microplastic pieces per <u>oyster</u> and nine per <u>clam</u> in the samples, and nearly all were microfibers, which can come from clothing made from synthetic or <u>natural materials</u> as well as derelict fishing gear.

"These microfilaments can be shed from clothing, up to 700,000 per load of laundry," Baechler said. "Those particles then travel out through greywater into wastewater and to the coast."

Microplastics were found in both Pacific oysters and razor clams collected from all 15 sample sites along the Oregon coast during spring and summer 2017. Microplastics were found in all but two of the roughly 300 organisms sampled.

The team also found that spring oysters contained more microplastics than the oysters sampled from the summer. Baechler said seasonal precipitation and the type of clothing typically worn in the spring compared to summer may have been factors.

"Whether it was a fairly urban site or a rural site, estuary or open-coast beach, both species had microplastics," Granek said. "Although we think of the Oregon coast as a much more pristine coastline compared to



California, Puget Sound or the Eastern Seaboard, when we are talking about microplastics, we're still seeing that human footprint on even our more pristine coastline."

Granek said that because fishing gear can be a source of microfibers, fisheries and oyster growers are often blamed for the problem of microplastics in seafood—but there is no scientific consensus that this is the case.

"It's not because people aren't managing our fisheries well or are being unclean in their practices," she said. "We're all using plastics on a daily basis. We are all the source of contamination in our seafood. And microplastics are not just in our seafood. We know that they are in our beer, in our salt, in our <u>drinking water</u>."

Baechler and Granek said more research still needs to be done to determine what effect the microplastics have on the organisms themselves and the humans who consume them. Studies have shown that microplastics can have negative physiological impacts such as reproductive and growth impairments on oysters and clams.

"If reproduction or growth is impaired, that could really affect not just individual clams or oysters, but possibly local populations of these organisms as well," Baechler said.

Granek said engineers are coming up with filters that could attach to washing machines, but it's still too early to tell how effective they are at preventing microfibers from discharging into the water and whether they're too costly for the general public.

The team's research was supported by Oregon Sea Grant, and their findings were published in the journal *Limnology and Oceanography Letters*.



More information: *Limnology and Oceanography Letters*, <u>DOI:</u> <u>10.1002/lol2.10124</u>

Provided by Portland State University

Citation: Microplastics found in oysters, clams on Oregon coast, study finds (2019, November 12) retrieved 2 May 2024 from https://phys.org/news/2019-11-microplastics-oysters-clams-oregon-coast.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.