

Millions of seabirds rely on discarded fish

November 8 2019



Credit: CC0 Public Domain

Millions of scavenging seabirds survive on fish discarded by North Sea fishing vessels, new research shows.

University of Exeter scientists estimate that 267,000 tonnes of fish was discarded in the North Sea in 2010—enough to feed 3.45 million birds.

This discard figure is down from almost 510,000 tonnes—enough for an

estimated 5.66 million birds—in 1990.

Discarding in the North Sea—one of the places in the world with the highest levels of this—is thought to have peaked around 1990.

The study examined eight [species](#), including [northern gannets](#) and herring gulls, and the figures are based on birds that rely to some extent on discarded fish (based on observations of how much discarded fish different bird species eat).

"Commercial fishing has a variety of effects on [marine life](#), but the impact of discards is one of the least studied and least understood," said lead author Dr. Richard Sherley, of the Environment and Sustainability Institute on Exeter's Penryn Campus in Cornwall.

"Our study highlights the sheer number of scavenging birds potentially supported by discards and thus the importance of understanding the wider ecological consequences of dumping fisheries waste.

"With discards declining over the period we studied, the number of [birds](#) able to rely on this has also declined."

The researchers estimate that the largest declines were in northern fulmars (1.4 million), black-legged kittiwakes (1.3 million) and herring gulls (630,000).

These declines also coincide with population declines at some North Sea colonies in each of these species.

However, the reasons for these declines are not entirely clear, and may not necessarily be underpinned by changes in discards, though changes in herring [gull](#) numbers at some sites have been linked to declining discards.

Dr. Sherley said fish are discarded for a variety of reasons.

"Fishers trying to catch one species may catch another (known as bycatch), and some fish may be partly processed on [fishing boats](#), with the resulting offal thrown overboard," he said.

"Changes including the EU's 'landing obligation' – which says any fish caught should be brought to shore—have reduced the amount of discard.

"There have also been improvements in fishery management, more targeted fishing—aided by technology to prevent bycatch—and efforts to land and use any [fish](#) that are caught, even if they're not among the species that humans generally eat."

The study, supported by the University of Kiel (Germany), considered the following seabird species: northern fulmar, northern gannet, great skua, common gull, lesser black-backed gull, herring gull, great black-backed gull and black-legged kittiwake.

The University of Exeter researchers included masters student Hannah Ladd-Jones and undergraduate Olivia Stevenson.

The paper, published in the journal *Fish and Fisheries*, is entitled: "Scavenger communities and fisheries waste: North Sea discards support 3 million seabirds, 2 million fewer than in 1990."

More information: Richard B. Sherley et al. Scavenger communities and fisheries waste: North Sea discards support 3 million seabirds, 2 million fewer than in 1990, *Fish and Fisheries* (2019). [DOI: 10.1111/faf.12422](https://doi.org/10.1111/faf.12422)

Provided by University of Exeter

Citation: Millions of seabirds rely on discarded fish (2019, November 8) retrieved 20 April 2024 from <https://phys.org/news/2019-11-millions-seabirds-discarded-fish.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.