

Water could modulate the activity and selectivity of carbon dioxide reduction

November 15 2019



The schematic diagram of water effect on CO_2 hydrogenation over stepped Cu(211) surface. Credit: Science China Press

As an alternative to the depletion of fossil resources, the reduction of



 CO_2 emitted from fossil fuel combustion into valuable chemicals and fuel has drawn increasing attention. Due to the highly thermodynamic stability of CO_2 , it is still very challenging to find a sustainable and costefficient route to selectively convert inert CO_2 at a high conversion rate under mild conditions. The water could also be a significant factor governing the reaction mechanism. On the one hand, water is one product accompanied with CH_3OH . H_2O might trigger the autocatalysis to accelerate CO_2 activation and methanol production. On the other hand, Cu is also a highly active catalyst for water-gas shift (WGS) reaction and its reverse reaction (RWGS). The water is also possible to launch or to suppress the side reaction, which could influence the selectivity.

Very recently, Dr. Cao and Prof. P. Hu's group at East China University of Technology systematically analyzed the effect of surface <u>water</u> on each elementary step to disclose its role in CO_2 hydrogenation over the stepped Cu(211) surface. The influence of water on CO_2 hydrogenation was respectively discussed at the levels of the electronic structure, energy and kinetics. The mechanisms between in the absence and presence of water were quantitatively compared based on microkinetic simulation upon density functional theory (DFT) calculations.

It has been found that the autocatalysis of water could be in favor of the formation of COOH intermediate through the proton transfer. This accelerates the CO generation while does not improve the methanol synthesis, giving rise to the higher CO_2 conversion and the selectivity towards CO. In addition, the microkinetic analysis also sheds light on the fact that the too high initial partial pressure of water will thermodynamically inhibit the CO_2 conversion due to the too high coverage of OH. Hence, it is the first time to disclose how to modulate the activity and selectivity of CO_2 hydrogenation through changing the H₂O pressure.



More information: Xitong Sun et al. A first-principles microkinetic study on the hydrogenation of carbon dioxide over Cu(211) in the presence of water. *Science China Chemistry*. doi.org/10.1007/s11426-019-9639-0

Provided by Science China Press

Citation: Water could modulate the activity and selectivity of carbon dioxide reduction (2019, November 15) retrieved 3 May 2024 from <u>https://phys.org/news/2019-11-modulate-carbon-dioxide-reduction.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.