

New study sheds more light on the properties of three polars

November 4 2019, by Tomasz Nowakowski



Optical spectrum of RX J0859.1+0537. Credit: Joshi et al., 2019.

Cataclysmic variables (CVs) are binary star systems consisting of a white



dwarf and a normal star companion. They irregularly increase in brightness by a large factor, then drop back down to a quiescent state. Although over 140 polars have been detected to date, only 33 of them have been identified as eclipsing systems. Observations of these rare objects could offer astronomers more opportunities to study magnetic accretion in binaries, for instance.

Indian astronomers have conducted an optical and x-ray study of three polars, a subclass of cataclysmic variables distinguished from other CVs by the presence of a very strong magnetic field in their white dwarfs. Specifically, they studied RX J0859.1+0537, RX J0749.1-0549 and RX J0649.8-0737. Results of the new study, presented in a paper published November 4 on arXiv.org, provide more details regarding fundamental properties of the three polars, which could advance the knowledge about such objects in general.

RX J0859.1+0537 (or RX J0859 for short), RX J0749.1-0549 (RX J0749), and RX J0649.8-0737 (RX J0649) are poorly studied polars located some 1,437 light years, 3,807 light years and 1,991 <u>light years</u> away from the Earth, respectively. In order to get more insights into the nature of the three polars, a team of astronomers led by Arti Joshi of Aryabhatta Research Institute of Observational Sciences in India performed optical photometric and spectroscopic observations of these objects. The study was complemented by an analysis of archival X-ray data.

"The poor knowledge of these sources encouraged us to do the optical (photometric and spectroscopic) and X-ray study," the astronomers wrote in the paper.

In general, the study confirmed that RX J0859, RX J0749 and RX J0649 belong to a class of eclipsing polars. The light curves of these sources showcase eclipse features that are deep, total and variable in shape. The



optical spectra of all the three systems exhibit strong, high-ionization emission lines and an inverted Balmer decrement, which confirms their magnetic nature.

According to the research, RX J0859, RX J0749 and RX J0649 have <u>orbital periods</u> of 2.39, 3.67 and 4.35 hours, respectively. Taking into account these results, the researchers classified RX J0859 as an eclipsing polar lying in the region of the period gap, while the other two were found to be long-period eclipsing polars above the period gap.

The astronomers found that for all the three systems, the radius of the eclipsed light source is found to be more than the actual size of the white dwarf. This suggest that the eclipsed component is not only a white dwarf, but may also be an extended accretion stream or an extended magnetic region on the white dwarf surface.

Moreover, the researchers identified cyclotron humps in the <u>optical</u> <u>spectra</u> of the three polars. This allowed them to measure the magnetic field strength of the surface of <u>white dwarfs</u> in these systems. This value was calculated to be at a level of about 49, 43.5, and 44 MG for RX J0859, RX J0749, and RX J0649, respectively.

More information: Optical and X-ray studies of three polars: RX J0859.1+0537, RX J0749.1-0549, and RX J0649.8-0737, arXiv:1910.11058 [astro-ph.SR] <u>arxiv.org/abs/1910.11058</u>

© 2019 Science X Network

Citation: New study sheds more light on the properties of three polars (2019, November 4) retrieved 23 June 2024 from <u>https://phys.org/news/2019-11-properties-polars.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.