


Scientists uncover resistance genes for deadly ash tree disease

November 18 2019



Credit: CC0 Public Domain

New research has identified the genetic basis of resistance to ash

dieback in UK trees, opening up new avenues for conservation.

Researchers from Queen Mary University of London and the Royal Botanic Gardens, Kew sequenced the DNA from over 1,250 [ash trees](#) to find inherited [genes](#) associated with ash [dieback](#) resistance.

The study, published in leading journal *Nature Ecology & Evolution*, showed that resistance is controlled by multiple genes, offering hope that surviving [trees](#) could be used to restore diseased woodlands, either by natural regeneration or [selective breeding](#).

Professor Richard Nichols, author of the study from Queen Mary University of London, said:

"We found that the genetics behind ash dieback resistance resembled other characteristics like human height, where the trait is controlled by many different genes working together, rather than one specific gene."

"Now we have established which genes are important for resistance we can predict which trees will survive ash dieback. This will help identify susceptible trees that need to be removed from woodlands, and provide the foundations for breeding more resistant trees in future."

Samples were collected from ash trees in a Forest Research mass screening trial, which comprises 150,000 trees planted across 14 sites in South East England.

The researchers screened for resistance genes using a rapid, cost-effective approach, where the DNA of multiple trees was combined into separate pools for diseased and unaffected trees.

Many of the genes found to be associated with ash dieback resistance were similar to those previously shown to be involved in disease or

pathogen responses in other species.

Ash dieback is a major threat to the UK landscape, and it is predicted it will kill over half of ash trees across the UK.

The disease, caused by the invasive alien fungus *Hymenoscyphus fraxineus*, has spread throughout Europe's ash populations, and was first recorded in the UK in 2012. It is estimated to cost the British economy £7.6 billion over the next 10 years.

Professor Richard Buggs, Senior Research Leader in Plant Health at the Royal Botanic Gardens, Kew and lead author of the paper, says:

"There is no cure for ash dieback and it threatens to kill over half of the 90 million ash trees in the UK. This will have huge impacts on the British landscape. Our new findings of the genetic basis of natural resistance found in a small minority of British ash trees help us to predict how ash populations will evolve under ash dieback. While many ash trees will die, our findings are encouraging from a long-term perspective and reassure us that ash woodlands will one day flourish again."

More information: Genomic basis of European ash tree resistance to ash dieback fungus, *Nature Ecology & Evolution* (2019). [DOI: 10.1038/s41559-019-1036-6](https://doi.org/10.1038/s41559-019-1036-6) , [nature.com/articles/s41559-019-1036-6](https://www.nature.com/articles/s41559-019-1036-6)

Tim L. R. Coker et al. Estimating mortality rates of European ash (*Fraxinus excelsior*) under the ash dieback (*Hymenoscyphus fraxineus*) epidemic, *PLANTS, PEOPLE, PLANET* (2018). [DOI: 10.1002/ppp3.11](https://doi.org/10.1002/ppp3.11)

Louise Hill et al. The £15 billion cost of ash dieback in Britain, *Current Biology* (2019). [DOI: 10.1016/j.cub.2019.03.033](https://doi.org/10.1016/j.cub.2019.03.033)

Provided by Queen Mary, University of London

Citation: Scientists uncover resistance genes for deadly ash tree disease (2019, November 18)
retrieved 19 April 2024 from

<https://phys.org/news/2019-11-scientists-uncover-resistance-genes-deadly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.