

## Better anchor roots help crops grow in poor soils

December 30 2019



A newly identified metabolite that stimulates anchor root growth in Arabidopsis plants may have potential applications in promoting plant growth in nutrient-deficient soils. The two plants treated with anchorene (right) show anchor root formation while untreated plants do not show anchor root formation. Credit: 2019 KAUST

A metabolite in plants that regulates the growth of anchor roots—vital for sustaining water and nutrient uptake in plants—has been identified and may have useful applications in agriculture.



Pigment compounds called carotenoids are found in all <u>plants</u> and play a key role in successful photosynthesis and the generation of plant hormones and metabolites. These products are formed when enzyme activity causes carotenoid molecules to split—a process known as cleavage. While many carotenoid products are known to play key biological roles, less is known about one group of cleavage molecules called di-apocarotenoids.

"Di-apocarotenoids have rarely been characterized due to their instability and low abundance," says KAUST research scientist Kunpeng Jia, who worked on the project under the supervision of KAUST's Salim Al-Babili. "Indeed, we are only beginning to understand what their biological significance might be and what functions they have."

The KAUST researchers, in collaboration with scientists in the United States and Germany, conducted an extensive study on the presence and biological activities of di-apocarotenoids in Arabidopsis plants using developmental studies and state-of-the-art analytical chemistry techniques. Working with such inherently unstable compounds that have low molecular weights within <u>plant tissues</u> was a real challenge for Jianing Mi, from Al-Babili's team, who honed lab techniques to extract and analyze the molecules without damaging them.

"We identified the di-apocarotenoid anchorene as a metabolite that sends a specific signal to trigger the formation of Arabidopsis anchor roots," says Jia. "Because anchorene is a carotenoid product, correct carotenoid biosynthesis is also required for healthy root formation. We confirmed this using chemical inhibitors and Arabidopsis mutants."

Further experiments showed that anchorene modulates the distribution of the plant hormone auxin in the anchor root formation site, which stimulates growth. Jia and coworkers found that increasing anchorene levels in <u>carotenoid</u>-deficient plants rescued anchor root growth, while



promoting growth in normal seedlings. When they modified anchorene's structure, it resulted in loss of activity.

"We'd like to explore the biological importance of anchorene further, and we also hope to understand exactly how plants produce this metabolite," says Jia. "We will also examine the biological activity of anchorene in crop plants because our findings may be relevant in boosting yields."

"Anchorene changes the root architecture by promoting anchor root formation, which increases root volume and facilitates water and nutrient absorption," adds Al-Babili. "Therefore, it may be possible to apply anchorene in nutrient-deficient soils to promote root growth."

**More information:** Kun-Peng Jia et al, Anchorene is a carotenoidderived regulatory metabolite required for anchor root formation in Arabidopsis, *Science Advances* (2019). <u>DOI: 10.1126/sciadv.aaw6787</u>

## Provided by King Abdullah University of Science and Technology

Citation: Better anchor roots help crops grow in poor soils (2019, December 30) retrieved 3 May 2024 from <u>https://phys.org/news/2019-12-anchor-roots-crops-poor-soils.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.