

Conferring leaf rust resistance in cereal crops

December 5 2019



Leaf rust infects wheat plants, considerably reducing crop yield. Credit: KAUST

Genes have been identified that confer resistance to multiple leaf rust species in barley. The findings by an international team, led by KAUST researchers, could transform the breeding of durable disease-resistant cereal crops and help support efforts to improve global food security.

"Disease is the exception and resistance is the rule—most microbes do not make us or cereal plants sick," says Simon Krattinger from



KAUST'sCenter for Desert Agriculture. "This is called nonhost resistance—resistance of an entire species against all strains of a pathogen. However, nonhost resistance in cereals is poorly understood."

The cereal-<u>rust</u> relationship is ideal for studying nonhost resistance because all cereals belong to the <u>grass family</u>, but each <u>cereal</u> crop species is infected by only one specific rust (for example, <u>wheat leaf</u> rust only infects wheat). There are molecular factors in barley that prevent wheat leaf rust from establishing colonies; thus, pinpointing the genes responsible for generating this molecular barrier to infection would be invaluable for breeders.

"Importantly, nonhost resistance is more durable than host resistance—a plant's innate immune system provides some protection, but only until pathogens evolve to evade it," says Krattinger's postdoc Yajun Wang. "Our biggest challenge was to identify the nonhost resistance genes in barley plants, especially given that barley's genome is almost twice the size of the human genome."

All barley cultivars are resistant to leaf rusts of other cereals; therefore, there is no clear genetic variation within barley species that might indicate which genes are involved. KAUST's collaborators in the Netherlands devised a novel method of narrowing the search.

They infected 1733 barley cultivars with wheat leaf rust. Most plants were resistant, but a few lines developed hints of leaf rust at the seedling stage. This was not enough to create a full infection, but the team was able to crossbreed these lines to generate one line that was highly susceptible to wheat leaf rust. This was then crossed with a normal barley cultivar and analyzed to pinpoint the genetic variations conferring nonhost resistance.





Wheat seeds are planted for trials in the KAUST Core Lab's Plant Growth Facility. Credit: KAUST

"Through extensive genome analysis, we found the genes that encode a protein receptor kinase to create a barrier to wheat rust in <u>barley</u>," says Wang. "Transferring these genes into wheat could result in cultivars that are resistant to all races of wheat rust."

"This is a very promising strategy that could finally solve one of the biggest problems in global wheat production," notes Krattinger.

More information: Wang, Y., et al. Orthologous receptor kinases quantitatively affect the host status of barley to leaf rust fungi. *Nature Plants* 5 1129–1135 (2019).



Provided by King Abdullah University of Science and Technology

Citation: Conferring leaf rust resistance in cereal crops (2019, December 5) retrieved 30 April 2024 from <u>https://phys.org/news/2019-12-conferring-leaf-rust-resistance-cereal.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.