


Finding your way in the dark depends on your internal clock

December 19 2019



Mouse in a maze in near darkness looking for a faint light which indicates the

exit. Credit: Ala-Laurila Lab

How mammals perceive light changes between night and day. Researchers at Aalto University and the University of Helsinki discovered that mice were better at finding a dim light in pitch-darkness in experiments done at night compared to those conducted during the day. The scientists were surprised to find that this effect had very little to do with any changes in the eye itself, and was instead controlled by how the mice actually searched for light in the dark, and how their brains processed signals at night vs. day. The results are exciting for neuroscientists interested in how animals and humans can see, and biologists interested in how the time of day alters our bodies, including sensory processing.

Many types of animals behave differently during day and night. While the body clock has long been known, its effect on body function still aren't fully understood. For instance, researchers knew that the retina, the part of the eye that detects light, has its own circadian rhythm. The team at Aalto and Helsinki Universities were interested in seeing if the eye's internal clock affected vision, so they modified one of their previous experiments to find out.

"Our research group is able to link if a mouse can find a dim light in the dark to the mouse's underlying retinal nerve signals at the sensitivity limit of vision," said Professor Petri Ala-Laurila, the research group leader. "This allowed us to explore how the day/night cycle changes the mouse's visual capability, both down in the neural circuit level and all the way up to behavior responses at the sensitivity limit of vision"

Finding your way in a pitch-dark maze

Earlier this year, [Professor Ala-Laurila's group demonstrated](#) how the eyes of mice detect faint light in near-total darkness. This allowed them to link the mammalian visually guided behavior to individual neural impulses, an important world-first for neuroscience. The experiment involved placing a mouse in a maze in total darkness with a faint light next to the exit of the maze. The mouse is trained to know that the light leads to the way out. They repeated this experiment for this new study, doing some of the tests during the day, and some during the night. They observed a behavior change—the mice were better at spotting the light at night than during the day. They were expecting this result, but they also observed that the nerve impulses leaving from the retina themselves did not cause the difference, which was a surprise.

So if the nerve signals from the eyes aren't changing, how could the mice be seeing better at night? The researchers were able to answer this using their new laboratory set-up. A big part of the technique invented by the group at Aalto involves using high-tech night vision cameras and their own deep-learning based software to track how the animals moved and what they could see. The team observed that during the night experiments, the mice searched for the light more effectively by scanning the environment, for example, by turning around more. Once the night group learned this behavioral strategy after searching at night, they were then also able to use it during the day, solving the maze puzzles quicker in day experiments than identical mice who'd never attempted it at night.

"Previously, it had not been possible to measure behavior as accurately as our group now can, so researchers had to treat [mice](#) as having a predefined set of behavioral rules in experiments like this. It's exciting to now show that even in the simplest of tasks—finding a light in the dark—animals can use vastly different behavioral strategies, and what's more, we are able to quantify day/night differences in them," said Sanna Koskela, a Ph.D. student at Helsinki University and the first author of

the paper in *Current Biology*, which published the results.

Internal Clock effects on the eye

The team now hope to further investigate the effects of the circadian rhythm on the eye. Although this specific test doesn't appear to show any signal effects from time of day, it is just one of many visual tasks the eye can perform at low light level, and others may yet still show circadian influence.

"We now have a remarkable opportunity to study sensory performance from the retina to behavior in [dim light](#), including things like how circadian rhythm controls it. Our next set of experiments will explore how the brain processes weak signals originating from increasing and decreasing light intensities in the retina at different times of day and night. This will help us understand more deeply how mammals see at low [light](#) levels," said Professor Ala-Laurila

More information: Mice reach higher visual sensitivity at night by using a more efficient behavioural strategy, *Current Biology*, [DOI: 10.1016/j.cub.2019.11.021](#)

Provided by Aalto University

Citation: Finding your way in the dark depends on your internal clock (2019, December 19) retrieved 8 April 2024 from <https://phys.org/news/2019-12-dark-internal-clock.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
