

First field measurements of laughing gas isotopes

December 5 2019



Empa researcher Erkan Ibraim controls one of the flux chambers that collect nitrous oxide emissions from the ground. Picture: Empa

Thanks to a newly developed laser spectrometer, Empa researchers can for the first time show which processes in grassland lead to nitrous oxide emissions. The aim is to reduce emissions of this potent greenhouse gas by gaining a better understanding of the processes taking place in the soil.

Nitrous oxide (N_2O , also known as laughing gas) is one of the most important greenhouse gases. Although it is much less abundant in the atmosphere than carbon dioxide (CO_2), it is around 300 times more potent. N_2O remains in the atmosphere for more than 100 years and thus contributes largely to global warming. It also damages the Earth's protective ozone layer. The largest source of N_2O emissions is soil—especially (but not only) when fertilized.

Researchers around the world are looking for ways to reduce N_2O emissions. But research is still in its infancy. "It is well known that more [nitrous oxide](#) escapes from the soil after fertilization or rainfall, for example. But little research has yet been done into the exact processes that take place in the soil," says Empa emissions and isotope researcher Joachim Mohn.

First measurements on grassland

Empa researchers have, therefore, developed a laser spectrometer, which enables extremely precise field measurements. "You can see exactly what isotope composition the emitted nitrous oxide has. For example, whether the [nitrogen atom](#) with an additional neutron is located in the middle of the molecule or at the edge," explains Mohn. The specific determination of the [isotopes](#) allows conclusions to be drawn about the formation processes of N_2O . "Isotope measurements can also be used to estimate the extent, to which the harmful nitrous oxide in the soil is degraded to harmless nitrogen."

N_2O is formed through various microbial processes. It can occur as a by-product of nitrification and as an intermediate product of denitrification. In nitrification, ammonium, e.g. from fertilizers, is oxidized to nitrate. In denitrification, nitrate is converted into nitrogen.

"Empa and other research institutions are currently investigating which biochemical process in a bacterium prefers to form which nitrous oxide isotope," says Mohn (see box). Based on these findings, Empa researchers, together with scientists from ETH Zurich and the Karlsruhe Institute of Technology (KIT), carried out more than 600 laser spectrometer measurements over several months in Bavaria over grassland and thus analyzed the isotope composition of the emitted N_2O .

At the same time, the researchers recorded influencing variables such as soil moisture, nutrient content, air temperature, wind speeds and the time of precipitation and fertilization. A novelty, as Joachim Mohn explains: "With the mass spectrometry instruments used so far it was simply impossible to measure continuously over [soil](#). Thanks to our new device, we can now carry out highly precise measurements in the field and compare the results, for example from grassland, with those from the lab."

The researchers are now using the first field measurements to check whether previous emission models allow good predictions or whether they need to be improved. Mohn: "So far, it has only been possible to say whether a model for predicting [nitrous oxide emissions](#) correctly reflects the time and quantity. If we also determine the isotope signature, then we know immediately whether the model correctly predicts the processes by which nitrous oxide is produced."

This is an important step for N_2O research, says the Empa researcher. "The long-term goal is to reduce nitrous [oxide](#) emissions from natural and agricultural soils." There is still a long way to go, he concedes. "But

at least we have now reached a first milestone."

More information: Attribution of N₂O sources in a grassland soil with laserspectroscopy based isotopocule analysis, *Biogeosciences*, doi.org/10.5194/bg-16-3247-201

Provided by Swiss Federal Laboratories for Materials Science and Technology

Citation: First field measurements of laughing gas isotopes (2019, December 5) retrieved 10 April 2024 from <https://phys.org/news/2019-12-field-gas-isotopes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.