

# Gene expression regulation in Chinese cabbage illuminated

December 5 2019




Figure 1: Tri-methylation of H3 lysine residue 27K (H3K27me3). Credit: Kobe University

Doctoral student Ayasha Akter (Kobe University's Graduate School of Agricultural Science) and technical staff member Satoshi Takahashi (from the RIKEN Center for Sustainable Resource Science) have revealed the important role played by the histone modification H3K27me3 in regulating gene expression in Chinese cabbage. In

addition, they illuminated the role of H3K27me3 in vernalization—a vital process for enabling *Brassica rapa* vegetables, such as Chinese cabbage, to flower.

The collaborative research group consisted of members from Kobe University's Graduate School of Agriculture, the RIKEN Center for Sustainable Resource Science, and CSIRO Australia.

These results were initially published online in the journal *DNA Research* on October 17, 2019.

Histones are proteins that play a role in [gene regulation](#)- they also package DNA into structures called nucleosomes. A nucleosome is a section of DNA that is wrapped around a core of eight histone proteins- this core is known as a histone octamer (Figure 1). Histone methylation is a process whereby certain amino acids in a histone protein are modified by the addition of one, two or three methyl groups. Methylation of histones changes the packaging of the DNA and allows [transcription factors](#) or other proteins to access the DNA and turn the [genes](#) on or off. In this way, histones can change [gene expression](#) but not the DNA sequence. This is known as an epigenetic change. In the case of H3K27me3, the methylation of the H3 family histone targets the lysine residues (K27). H3K27me3 is often associated with gene silencing.




Figure 2: Graph to show the difference in the level of H3K27me3 in genes between the two lines and between tissues. T23 and T24 are the lines, and 2d-C and 14d-L are tissues. Credit: Kobe University

The whole genome sequence for Chinese cabbage (*Brassica rapa* L.) has been available since 2011. However, the exact role that H3K27me3 plays was not well understood. This study revealed the importance of H3K27me3 in vernalization—the process by which Chinese cabbage flowers after cold exposure.

For this study, two inbred lines of Chinese cabbage were used (T23 and T24). These lines were selected to investigate the role of H3K27me3 in

tissue-specific gene expression because they are genetically similar to the reference whole genome sequence that was discovered in 2011. The research team compared the number of genes marked by H3K27me3 between the two lines, and between different tissues. They found that the distribution of H3K27me3 was similar between the two lines. However, there was greater variation in H3K27me3 distribution across different tissues within a line. In other words, it was found that genes with H3K27me3 marks showed tissue-specific expression (Figure 2). This suggests that this histone modification plays a role in the regulation of gene expression during plant tissue development.

Next, they investigated the relationship between the level of gene expression and the presence of H3K27me3. To do this, they measured the level of expression in the genes that were modified by H3K27me3 in all the samples (around 8000 genes), and compared this with the level of expression in all the genes (around 40,000 genes). They found that the level of expression in genes that were modified by H3K27me3 was lower (Figure 3). These results revealed that H3K27me3 plays a role in repressing gene expression. Furthermore, it was found that the level of H3K27me3 was high in genes responsible for tissue-specific expression (Figure 4). This suggests that H3K27me3 regulates gene expression during plant development.




Figure 3: The expression levels of genes with H3K27me3. 'K27+' indicates genes with H3K27me3, and 'Total' refers to all the genes. Credit: Kobe University

Chinese cabbage is from the same *Brassica* genus of plants as *Arabidopsis thaliana*. Therefore, the researchers decided to compare the presence of H3K27me3 in these closely related plants. It was found that 40% of the modified genes in the two species were altered in a similar way by H3K27me3, suggesting that these modifications remained after speciation. This showed that H3K27me3 modification plays a vital role in both species.

Lastly, an experiment was conducted to see how H3K27me3 levels

changed before and after vernalization. Vernalization refers to the process in which some varieties of plant flower after a period of exposure to cold weather. In a previous study it was revealed that the gene FLOWERING LOCUS C (FLC) plays a key role in preventing vegetables in the *B. rapa* family from flowering before vernalization. The expression of the FLC gene ensures that the plant cannot flower before it is exposed to cold weather- this helps prevent the plant from flowering in winter. The FLC gene is repressed when the plant undergoes a cold period- allowing it to flower once temperatures become warmer.

To mimic the conditions for vernalization, Chinese cabbage and *Arabidopsis thaliana* plants were kept at low temperatures of 4°C for a 4 week period. The results found that there were no large changes in H3K27me3 levels at a genome level. However, H3K27me3 accumulated in the FLC genes of both species after cold exposure. In Chinese cabbage, it was observed that H3K27me3 accumulated near FLC transcription initiation sites. In other words, H3K27me3 levels were high where the FLC genes were turned 'off.' When normal growing temperatures returned (22°C), H3K27me3 spread along the entirety of the FLC paralogs. This suggests that H3K27me3 also plays an important role in ensuring that the FLC gene remains repressed after vernalization.




Figure 4: Tissue-specific expression in genes with H3K27me3. 'K27+' indicates genes with H3K27me3, and 'Total' refers to all the genes. Credit: Kobe University




Figure 5: Accumulation of H3K27me3 in FLC before and after a period of cold exposure. The graph shows the results for BrFLC1. The same results were found in the paralogs BrFLC2, BrFLC3 and BrFLC5. Credit: Kobe University

Overall, this research illuminated that the histone modification H3 lysine 27 tri-methylation plays important roles in regulating the expression of genes responsible for tissue differentiation. They also revealed the indispensable role of H3K27me3 in controlling the expression of FLC during vernalization.

Further research into the role H3K27me3 plays in individual genes, as well as into any differences between H3K27me3 levels in different cultivars would deepen understanding around the role of this histone modification.

If successful, experiments in using H3K27me3 to suppress FLC could enable flowering time to be controlled artificially. This knowledge could be utilized in agriculture to improve the cultivation rates of vegetables in the *B. rapa* family.

**More information:** Ayasha Akter et al. The histone modification H3 lysine 27 tri-methylation has conserved gene regulatory roles in the triplicated genome of *Brassica rapa* L., *DNA Research* (2019). [DOI: 10.1093/dnares/dsz021](https://doi.org/10.1093/dnares/dsz021)

Provided by Kobe University

Citation: Gene expression regulation in Chinese cabbage illuminated (2019, December 5) retrieved 16 June 2024 from <https://phys.org/news/2019-12-gene-chinese-cabbage->



[illuminated.html](http://illuminated.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.