

New rules illuminate how objects absorb and emit light

December 23 2019, by Scott Lyon



Princeton researchers, led by Alejandro Rodriguez, have uncovered new rules for how objects absorb and emit light. The work resolves a longstanding discrepancy between large and small objects, unifying the theory of thermal radiation across all scales and boosting scientists' control in designing light-based technology. Credit: Casey Horner/Unsplash

Princeton researchers have uncovered new rules governing how objects absorb and emit light, fine-tuning scientists' control over light and boosting research into next-generation solar and optical devices.

The discovery solves a longstanding problem of scale, where light's behavior when interacting with [tiny objects](#) violates well-established physical constraints observed at larger scales.

"The kinds of effects you get for very small objects are different from

the effects you get from very large objects," said Sean Molesky, a postdoctoral researcher in electrical engineering and the study's first author. The difference can be observed in moving from a molecule to a grain of sand. "You can't simultaneously describe both things," he said.

The problem stems from light's famous shapeshifting nature. For ordinary objects, light's movement can be described by straight lines, or rays. But for microscopic objects, light's wave properties take over and the neat rules of ray optics break down. The effects are significant. In important modern materials, observations at the micron scale showed [infrared light](#) radiating at millions of times more energy per unit area than ray optics predicts.

The new rules, published in *Physical Review Letters* on Dec. 20, tell scientists how much infrared light an [object](#) of any scale can be expected to absorb or emit, resolving a decades-old discrepancy between big and small. The work extends a 19th-century concept, known as a blackbody, into a useful modern context. Blackbodies are idealized objects that absorb and emit light with maximum efficiency.

"There's been a lot of research done to try to understand in practice, for a given material, how one can approach these blackbody limits," said Alejandro Rodriguez, an associate professor of electrical engineering and the study's principal investigator. "How can we make a perfect absorber? A perfect emitter?"

"It's a very old problem that many physicists—including Planck, Einstein and Boltzmann—tackled early on and laid the foundations for the development of quantum mechanics."

A large body of previous work has shown that structuring objects with nanoscale features can enhance absorption and emission, effectively trapping photons in a tiny hall of mirrors. But no one had defined the

fundamental limits of the possible, leaving open major questions about how to assess a design.

No longer confined to brute-force trial and error, the new level of control will allow engineers to optimize designs mathematically for a wide range of future applications. The work is especially important in technologies like solar panels, optical circuits and quantum computers.

Currently, the team's findings are specific to thermal sources of light, like the sun or like an incandescent bulb. But the researchers hope to generalize the work further to agree with other [light](#) sources, like LEDs, fireflies, or arcing bolts of electricity.

More information: Sean Molesky et al, T Operator Bounds on Angle-Integrated Absorption and Thermal Radiation for Arbitrary Objects, *Physical Review Letters* (2019). [DOI: 10.1103/PhysRevLett.123.257401](https://doi.org/10.1103/PhysRevLett.123.257401)

Provided by Princeton University

Citation: New rules illuminate how objects absorb and emit light (2019, December 23) retrieved 2 May 2024 from <https://phys.org/news/2019-12-illuminate-absorb-emit.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--