

Scientists use air-based maps of plant chemistry to improve carbon cycling models in hyperdiverse tropical forests

December 4 2019



Images of the lowland Amazon rainforest canopy. Different colors represent different spectral fingerprints. Credit: Global Airborne Observatory (GAO)

An international team of scientists led by the University of Arizona used the latest technology in remote sensing to measure plant biodiversity from the Amazon basin to the Andes Mountains in Peru to better



understand how tropical forests will respond to climate change.

The researchers used Arizona State University's Global Airborne Observatory, or GAO, to show that by combining traditional on-theground field measurements of carbon with aerial measurements of plant chemistry, the ability to model and predict the role that tropical forests play in the <u>global carbon cycle</u> can be improved.

"This work is important because it can be difficult to obtain measurements in some of these remote places," said Sandra Durán, a postdoctoral fellow at the University of Arizona and lead author of the paper published in *Science Advances*.

"Climate scientists are interested in predicting how much carbon is able to be captured by specific forests. We're showing that these measurements of plant chemistry taken from airplanes have potential to make predictions of carbon gain in one of the most biodiverse forests in the world for the first time."

A tree's ability to grow and survive is affected by traits such as nutrient concentrations and defense compounds in its leaves. While researchers have measured these traits in trees located around the globe, incomplete data from the Andes' highly diverse forests has made it difficult to understand how trees impact the functioning of tropical forests.

Durán and her collaborators utilized GAO maps of plant chemistry to quantify the diversity of plant chemistry that is not visible to the naked eye. The measurements allowed them to see how diversity can predict rates of carbon capture in tropical forests that contain a large range of temperatures and elevations.

"New technology is enabling us to see the functioning of the <u>forest</u> in a new light," said corresponding author Brian Enquist, professor of



ecology and evolutionary biology at the University of Arizona. "This technology creates a continuous map of the variation in plant chemistry, even in highly remote areas of the Earth that would be almost impossible to gather from ground surveys. This will improve our ability to model tree responses to environmental changes, from small plots to large regions."

Greg Asner, co-author and principal investigator of GAO, said, "Our plant canopy chemistry maps have been used for many purposes over the past 10 years, but this new application—to assess drivers of forest carbon cycling—is new and opens doors for the use of our mapping approach throughout the world's <u>tropical forests</u>."

More information: "Informing trait-based ecology by assessing remotely sensed functional diversity across a broad tropical temperature gradient" *Science Advances* (2019). DOI: 10.1126/sciadv.aaw8114, advances.sciencemag.org/content/5/12/eaaw8114

Provided by University of Arizona

Citation: Scientists use air-based maps of plant chemistry to improve carbon cycling models in hyperdiverse tropical forests (2019, December 4) retrieved 1 May 2024 from <u>https://phys.org/news/2019-12-scientists-air-based-chemistry-carbon-hyperdiverse.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.