

Uranium chemistry and geological disposal of radioactive waste

December 16 2019



Demonstration of a sample containing uranium being studied and carefully loaded onto Diamond's I20-scanning beamline - Credit Diamond Light Source
Credit: Diamond Light Source

A new paper to be published on 16 December provides a significant new insight into our understanding of uranium biogeochemistry and could help with the UK's nuclear legacy.


Conducted by a team of researchers from the University of Manchester, Diamond Light Source and Radioactive Waste Management, their work shows for the first time how uranium forms a uranium-sulfur complex under conditions generally found in the environment and how this compound can be an important intermediary in uranium immobilization. Published in *Environmental Science & Technology*, the paper is called "Formation of a U(VI)-persulfide complex during environmentally relevant sulfidation of iron (oxyhydr)oxides."

Professor Katherine Morris, associate dean for research facilities in the Faculty of Science and Engineering, University of Manchester and the research director for the BNFL Research Centre in Radwaste Disposal explains why recreating and studying these chemical complexes is highly relevant for understanding and dealing with radioactive waste: "To be able to predict the behavior of the uranium during geological disposal, we need to take into account that it may have interacted with other processes taking place in the ground. These so-called biogeochemical reactions are often a complex set of interactions between dissolved chemical species, mineral surfaces, and microorganisms."

The recent study is the first time that researchers have shown that a uranium-sulfide complex can form under conditions representative of a deep underground environment. This complex then transforms further into highly immobile uranium oxide nanoparticles.

In the experiment, the researchers studied uranium when it sits at the surface of the mineral ferrihydrite, which is a widespread mineral in the environment. The researchers used an X-ray based method called X-ray Absorption Spectroscopy (XAS) to study the samples at Diamond Light

Source, the UK's national Synchrotron. The XAS data, in combination with computational modeling, showed that during the sulfidation reaction, a short-lived and novel U(VI)-persulfide complex formed during this biogeochemical process.



The uranium-persulfide complex associated with the transforming mineral surface. Credit: Diamond Light Source

Professor Sam Shaw, co-investigator and professor of environmental mineralogy at the University of Manchester; "Shining the synchrotron beam onto the sample causes the uranium within to emit X-rays. By analyzing the X-ray signal from the samples our team were able to determine the chemical form of uranium, and to which other elements it is bound. To further validate the theory on the formation pathway of the uranium-sulfur complexes, our team also made computer simulations to conclude which type of complex is more likely to form. This is the first observation of this form of uranium under aqueous conditions, and provides new insight into how uranium behaves in environments where sulfide is present. This work demonstrates the deep understanding we can develop of these complex systems and this knowledge will help underpin efforts to manage radioactive wastes in a geological disposal facility."

Dr. Luke Townsend, postdoctoral fellow in environmental radiochemistry at the University of Manchester, who undertook this research as part of his Ph.D., further adds, "When trying to mimic environmental processes in the laboratory, it's a challenge to produce accurate, high quality, reproducible science with such complex experiments, whilst also maintaining relevance to the geodisposal environment. However, obtaining exciting results such as these makes all the hard work and commitment to the project from myself and the group, both in our labs in Manchester and on the beamlines at Diamond, completely worthwhile."

The XAS measurements were performed at Diamond on beamlines I20 and B18 by the researchers who used highly controlled sulfidation experiments that mimic biogeochemical processes in the deep underground environment. This was combined with geochemical analyses and computational modeling to track and understand uranium behavior.

Physical science director at Diamond, Laurent Chapon, concludes, "This is another example of how Diamond's state of the art analytical tools are enabling scientists to follow complex processes and help them to tackle 21st century challenges. In this instance, our beamlines allowed the users to gain real insight into the environmental relevance of this new [uranium](#)-sulfur complex, which feeds into our understanding of geological disposal."

More information: "Formation of a U(VI)-persulfide complex during environmentally relevant sulfidation of iron (oxyhydr)oxides" *Environmental Science & Technology* (2019). [DOI: 10.1021/acs.est.9b03180](#)

Provided by Diamond Light Source

Citation: Uranium chemistry and geological disposal of radioactive waste (2019, December 16) retrieved 27 June 2024 from <https://phys.org/news/2019-12-uranium-chemistry-geological-disposal-radioactive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.