

# Study finds deforestation is changing animal communication

January 9 2020

---



Credit: CC0 Public Domain

Deforestation is changing the way monkeys communicate in their natural habitat, according to a new study.

This study, led by an anthropologist at the University of Waterloo, offers the first evidence in animal communication scholarship of differences in vocal behaviours in response to different types of [forest](#) edge areas.

Working in a tropical lowland rainforest in Costa Rica, the researchers examined how human-caused forest habitat changes have affected vegetation and, in turn, the rate and length of howling by the group-living [howler monkey](#) species.

Led by Laura Bolt, an adjunct professor of anthropology at Waterloo, the study compared how the communication behaviour of the mantled howler monkey differs in forest edges impacted by [human activity](#), known as [anthropogenic](#) edges, compared to natural forest edges.

"Howler [monkeys](#) are well-known for making very loud, long-distance vocalizations called howls," said Bolt. "While howls are only produced by [adult males](#), howl function is not entirely known, so we conducted our study to test the hypothesis that the intensity of howling by monkeys relates to defending ecological resources such as areas of richer vegetation or preferred feeding trees."

Anthropogenic areas were identified as areas within 50 meters of barbed wire fences marking the edge of the forest and the start of coconut plantations or cattle pasture, and natural forest edges as areas within 50 meters of a river.

The study found that males howl to defend high-quality resources, with notably longer durations of howling in the forest interior and at river edge areas where vegetation resources are richer. The researchers also found differences in howl length between river edge and anthropogenic edge areas, which is an important insight for conservation planning.

"Howler monkeys eat leaves and fruit, and if they are howling to defend

these resources, we predicted that males would howl for longer durations of time when in a forest interior or near the river edge, where vegetation is richer compared to anthropogenic edge," said Bolt.

To conduct their study, the researchers collected data on mantled howler monkey howling behaviour from May to August in 2017 and 2018, following groups as they travelled across various edge and interior habitat zones. All monkey groups were well-habituated and did not react to the visible presence of the researchers.

With their evidence showing that anthropogenic deforestation is altering howler monkey behaviour, Bolt and her colleagues say that long-term howler monkey conservation initiatives should prioritize preservation of forest interior and river edge regions and re-forestation of human-caused forest edges.

"While it is yet unknown what implications these behavioural changes across different edge zones may have for monkey fitness," says Bolt, "our findings show that it is proximity to anthropogenic forest edge, rather than to naturally-occurring forest edge, that is changing howler monkey communication [behaviour](#). This is just one of the many ways that howler monkeys are affected by deforestation."

The study will be published in the journal *Behaviour* in spring 2020.

Provided by University of Waterloo

Citation: Study finds deforestation is changing animal communication (2020, January 9) retrieved 8 September 2024 from <https://phys.org/news/2020-01-deforestation-animal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.