


# Moving domain walls induce losses in superconductor/ferromagnet hybrid systems

January 6 2020



Josephson Junction. Credit: The University of Jyväskylä/ Mihail Silaev

Physicists have shown that the motion of domain walls can be detected by monitoring voltage generated in superconducting devices. This finding can facilitate magnetic racetrack memory applications. The result was published in *Physical Review Letters*. The international

research group included researchers from the University of Jyväskylä.

Recently many research groups have aimed at developing magnetic memories which are based on writing and reading magnetic information with the help of electric current. Such systems typically require such large amounts of current to switch magnetization that it can affect the thermal stability of the memory element. In order to reduce heating effects, [superconducting materials](#) which can sustain dissipationless electric current would be very useful.

Superconducting current is a flow of electrons bound in Cooper pairs, and therefore is fundamentally different from the usual current in normal metals which is carried by single electrons. Therefore, in order to develop a superconductor/ferromagnet [memory](#) element it has been necessary to understand how superconducting current can affect magnetic state.

In their paper, the research group has found answers to two fundamental questions: whether supercurrent can change magnetic states, and whether it is possible to avoid electric losses during this process.

"We have developed the theory which describes how the superconductor can lose its [fundamental property](#) of having zero resistance in a typical superconductor/ferromagnet device. This happens because of the induced magnetization dynamics in the attached ferromagnet. Although the force which drives magnetization comes from the superconducting current, the system becomes inherently dissipative and in principle cannot sustain any amount of superconducting current because of the voltage generated by the magnetization dynamics," explains Academy Research Fellow Mihail Silaev.

"We find the low-current resistance associated with the domain wall motion driven by the superconducting current. We suggest the finite

slope of Shapiro steps as the characteristic feature of the regime with domain wall oscillations driven by the external current flowing through the junction," Silaev concludes.

**More information:** D. S. Rabinovich et al, Resistive State of Superconductor-Ferromagnet-Superconductor Josephson Junctions in the Presence of Moving Domain Walls, *Physical Review Letters* (2019). [DOI: 10.1103/PhysRevLett.123.207001](https://doi.org/10.1103/PhysRevLett.123.207001)

Provided by University of Jyväskylä

Citation: Moving domain walls induce losses in superconductor/ferromagnet hybrid systems (2020, January 6) retrieved 8 September 2024 from <https://phys.org/news/2020-01-domain-walls-losses-superconductorferromagnet-hybrid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.