

## A quantum of solid: A glass nanoparticle in the quantum regime

January 31 2020



Scientists from Vienna, Kahan Dare (left) and Manuel Reisenbauer (right) working on the experiment that cooled a levitated nanoparticle to its motional quantum groundstate. Credit: © Lorenzo Magrini, Yuriy Coroli/University of Vienna

Researchers in Austria have used lasers to levitate and cool a glass



nanoparticle into the quantum regime. Although it is trapped in a roomtemperature environment, the particle's motion is solely governed by the laws of quantum physics. The team of scientists from the University of Vienna, the Austrian Academy of Sciences and the Massachusetts Institute of Technology (MIT) published their new study in the journal *Science*.

It is well known that quantum properties of individual atoms can be controlled and manipulated with <u>laser light</u>. Even large clouds of hundreds of millions of atoms can be pushed into the <u>quantum regime</u>, giving rise to macroscopic quantum states of matter such as quantum gases or Bose-Einstein condensates, which today are also widely used in quantum technologies. An exciting next step is to extend this level of quantum control to solid-state objects. In contrast to atomic clouds, the density of a solid is a billion times higher, and all atoms are bound to move together along the object's center of mass.

However, entering this new regime is not at all a straightforward endeavor. A first step for achieving such quantum control is to isolate the object under investigation from influences of the environment and to remove all thermal energy—by cooling it down to temperatures very close to absolute zero (-273.15 degrees Celsius) such that quantum mechanics dominates the particle's motion. To show this, the researchers chose to experiment with a glass bead approximately 1000 times smaller than a grain of sand and containing a few hundred million atoms. Isolation from the environment is achieved by optically trapping the particle in a tightly focused laser beam in high vacuum, a trick that was originally introduced by Nobel laureate Arthur Ashkin many decades ago, and that is also used for isolating atoms. "The real challenge is to cool the particle motion into its quantum ground state. Laser cooling via atomic transitions is well-established and a natural choice for atoms, but it does not work for solids," says lead author Uros Delic from the University of Vienna.



For this reason, the team has been working on implementing a lasercooling method that was proposed by Austrian physicist Helmut Ritsch at the University of Innsbruck and, independently, by study co-author Vladan Vuletic and Nobel laureate Steven Chu. They had recently announced a first demonstration of the working principle, cavity cooling by coherent scattering; however, they were still limited to operating far away from the quantum regime.

"We have upgraded our experiment and are now able not only to remove more background gas, but also to send in more photons for cooling," says Delic. In that way, the motion of the glass bead can be cooled straight into the quantum regime. "It is funny to think about this: The surface of our glass bead is extremely hot, around 300 degrees Celsius, because the laser heats up the electrons in the material. But the motion of the center of mass of the particle is ultra-cold, around 0.00001 degrees Celsius away from absolute zero, and we can show that the hot particle moves in a quantum way."





Researchers cooled a levitated nanoparticle to the quantum groundstate for the first time. This work was made possible by the recent breakthrough application of coherent scattering in the field of cavity optomechanics. Credit: Kahan Dare, Lorenzo Magrini, Yuriy Coroli, University of Vienna

The researchers are excited about the prospects of their work. The quantum motion of solids has also been investigated by other groups all around the world, along with the Vienna team. Thus far, experimental systems consisted of nano- and micromechanical resonators—in essence, drums or diving boards that are clamped to a rigid support structure. "Optical levitation brings in much more freedom: By changing the optical trap—or even switching it off—we can manipulate the nanoparticle motion in completely new ways," says Nikolai Kiesel, co-author and Assistant Professor at the University of Vienna.



Several schemes along these lines have been proposed, amongst others by Austrian-based physicists Oriol Romero-Isart and Peter Zoller at Innsbruck, and may now become possible. For example, in combination with the newly achieved motional ground state the authors expect that this opens new opportunities for unprecedented sensing performance, the study of fundamental processes of heat engines in the quantum regime, as well as the study of quantum phenomena involving large masses. "A decade ago, we started this experiment motivated by the prospect of a new category of quantum experiments. We finally have opened the door to this regime."

**More information:** U. Delić el al., "Cooling of a levitated nanoparticle to the motional quantum ground state," *Science* (2020). <u>science.sciencemag.org/lookup/ ... 1126/science.aba3993</u>

## Provided by University of Vienna

Citation: A quantum of solid: A glass nanoparticle in the quantum regime (2020, January 31) retrieved 1 May 2024 from https://phys.org/news/2020-01-quantum-solid-glass-nanoparticle-regime.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.