



Residues in fingerprints hold clues to their age

January 22 2020

Day 0



Day 1



Day 3



Levels of an unsaturated triacylglycerol decline in fingerprints from an individual from day 0 (top) to day 1 (middle) and day 3 (bottom). Credit: Adapted from *Analytical Chemistry* 2020, DOI: [10.1021/acs.analchem.9b04765](https://doi.org/10.1021/acs.analchem.9b04765)

Police have long relied on the unique whorls, loops or arches encoded in fingerprints to identify suspects. However, they have no way to tell how long ago those prints were left behind—information that could be crucial to a case. A preliminary new study in ACS' *Analytical Chemistry* suggests that could change. Researchers report that they could link compounds contained in fingerprints with their age.

By determining the age of [fingerprints](#), police could get an idea of who might have been present around the time a crime was committed. This information could, for example, contradict a suspect's explanation that he or she had visited earlier. Scientists have already begun mining fingerprint residues for clues to the identity of the person who made them, but timing has proven more difficult to reliably pin down. Notably, past research has shown that a gas chromatography-mass spectrometry method succeeded in determining if prints were more or less than eight days old; however, investigators often need more precision. To get a better idea of when prints were deposited, Young Jin Lee and colleagues looked to reactions already suspected to take place in these residues, when ozone in air reacts with unsaturated triacylglycerols left by a fingertip.

Using prints collected from three donors, the researchers tracked shifting levels of triacylglycerols using [mass spectrometry](#) imaging, a technique that leaves the evidence intact. They found they could reliably determine the triacylglycerol degradation rate for each person over the course of seven days. But the rate differed among individuals, with one person's triacylglycerols declining more gradually than the others. The researchers attribute this difference to higher levels of lipids in that individual's fingerprints. The method also worked on residues that had been dusted with forensic powder. The researchers say that although a large-scale study is needed to better understand how lipid levels affect

triacylglycerol degradation, this analysis is a first step toward developing a better fingerprint aging test.

More information: Paige Hinnert et al. Determining Fingerprint Age with Mass Spectrometry Imaging via Ozonolysis of Triacylglycerols, *Analytical Chemistry* (2020). [DOI: 10.1021/acs.analchem.9b04765](https://doi.org/10.1021/acs.analchem.9b04765)

Provided by American Chemical Society

Citation: Residues in fingerprints hold clues to their age (2020, January 22) retrieved 27 June 2024 from <https://phys.org/news/2020-01-residues-fingerprints-clues-age.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.