

## Smaller detection device effective for nuclear treaty verification, archaeology digs

January 29 2020



The schematic of a compact neutron resonance transmission analysis platform using a deuterium-tritium source (not to scale), shielding, a moderator for slowing down the neutrons to the eV regime and a boron-lined collimator. Credit: Ezra M. Engel and Ethan A. Klein, MIT

Most nuclear data measurements are performed at accelerators large enough to occupy a geologic formation a kilometer wide, like the Los Alamos Neutron Science Center located on a mesa in the desert. But a



portable device that can reveal the composition of materials quickly onsite would greatly benefit cases such as in archaeology and nuclear arms treaty verification.

Research published this week in *AIP Advances* used computational simulations to show that with the right geometric adjustments, it is possible to perform accurate neutron resonance transmission analysis in a <u>device</u> just 5 meters long.

"We expected massive backgrounds to dilute and contaminate our signal, and early simulation work confirmed that the scale of these effects would make the technique entirely impossible," author Areg Danagoulian said. "However, careful optimization of the geometries allowed us to almost completely suppress these effects, giving us a nearperfect signal."

Using a model of a pulsed deuterium-tritium neutron source in a polycone layout, the researchers performed a series of tests to optimize the moderation, shielding and collimation of the device and probe the configuration for uncertainties introduced by these adjustments. To confirm the device's accuracy, they compared spectral reconstructions and tested the isotopic sensitivity of the device.

"Depending on the goal of the application, one can use spectroscopic radiography to determine the absolute abundances and densities of individual isotopes," Danagoulian said. "It can also be used in treaty verification exercises, where an authentic nuclear weapon component is compared to that from a candidate warhead."

While the tests used silver, tungsten and molybdenum, the method could be used to identify isotopes of plutonium or uranium in nuclear warheads or enriched fuel, as well as tin, silver or gold in archaeological sites. Their work could also be used to similarly reduce the lengths of



thermal neutron beamlines.

Their work uses time-of-flight reconstructions of the energies of pulsed neutrons in order to determine the composition of target materials. These reconstructions allow analysis of the spectrum transmitted and nuclear resonances present in different isotopes to identify the isotopic makeup of the material in the target.

Their results show the device was successful. It was able to precisely differentiate various isotopes and was sensitive to variations in isotopic concentrations.

The authors plan to perform experimental validations of the above technique using various pulsed neutrons sources and <u>neutron</u> detectors.

**More information:** Ezra M. Engel et al, Feasibility study of a compact neutron resonance transmission analysis instrument, *AIP Advances* (2020). DOI: 10.1063/1.5129961

## Provided by American Institute of Physics

Citation: Smaller detection device effective for nuclear treaty verification, archaeology digs (2020, January 29) retrieved 1 May 2024 from <u>https://phys.org/news/2020-01-smaller-device-effective-nuclear-treaty.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.