


A better estimate of water-level rise in the Ganges delta

January 6 2020



A colourful view of the Bangladesh coastline seen by Envisat's Medium Resolution Imaging Spectrometer (MERIS). The image was acquired on 8 November 2003 in Full Resolution mode, providing a spatial resolution of 300 metres and covering approximately 633 km by 630 km. Credit: ESA, CC BY-SA 3.0 IGO

For the first time, scientists have provided reliable regional estimates of land subsidence and water-level rise in the Ganges-Brahmaputra-Meghna delta. Depending on the region of the delta, water-level rise could reach 85 to 140 cm by 2100. The work, published in *PNAS* on 6 January 2020 by researchers from the CNRS, IRD, BRGM, La Rochelle Université, Université des Antilles and Bangladesh University of Engineering and Technology¹, should provide input for future impact studies and adaptation plans.

Although the Ganges-Brahmaputra-Meghna delta is the largest and most densely populated delta in the world², and one of the places most vulnerable to [climate change](#), the extent and impact of water-level rise there remain poorly known. The area, which covers two thirds of Bangladesh and part of eastern India, is already regularly prone to flooding, favored by intense monsoon rainfall, rising sea levels, river flows and land [subsidence](#). However, it is difficult to disentangle these various factors. In addition, forecasts carried out so far have been based on highly localized measurements of water level.

To overcome these difficulties, the researchers analyzed monthly readings from 101 gauges measuring water and sea levels across the delta. By aggregating the data over geographical areas in order to filter out local effects and offset quality differences between gauges, they obtained robust estimates for water-level changes. Between 1968 and

2012, water level increased by 3 mm/year on average, slightly more than global mean [sea level](#) rise (2 mm/year during the same period).

They then estimated the contribution of land subsidence, obtained by subtracting from absolute sea level³ the measurements of relative water level obtained previously. According to their calculations, maximum land subsidence in the delta between 1993 and 2012 was between 1 and 7 mm/year. Although not negligible, these values are lower than some local measurements (e.g. 1-2 cm/year in Dhaka) that have been used as a reference until now.

If subsidence continues at the same rate, and even under a greenhouse gas mitigation scenario, water-level rise in the delta, depending on the area, could reach 85 to 140 cm by the end of the century compared to the period 1986-2005. This is twice as high as the projections provided in the latest IPCC report, which did not take land subsidence into account.

This work should help to improve forecasts of water level in the Ganges-Brahmaputra-Meghna [delta](#), and thus lead to refine impact studies and possible adaptation scenarios for its 200 million inhabitants.

More information: Mélanie Becker et al., "Water level changes, subsidence, and sea level rise in the Ganges–Brahmaputra–Meghna delta," *PNAS* (2019). www.pnas.org/cgi/doi/10.1073/pnas.1912921117

Provided by CNRS

Citation: A better estimate of water-level rise in the Ganges delta (2020, January 6) retrieved 3 May 2024 from <https://phys.org/news/2020-01-water-level-ganges-delta.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.