

Boost soybean yields by adapting photosynthesis to fleeting shadows, according to model

February 24 2020



University of Illinois graduate student Elsa de Becker (left) and postdoctoral researcher Steven Burgess (right) collect soybean leaf samples to use in a model that measures how well crops adapt to minute-by-minute changes in light intensity. They found that grappling with these fluctuations can cost 13% of the soybean crop's yield potential. Credit: Claire Benjamin/RIPE project

Komorebi is a Japanese word that describes how light filters through leaves—creating shifting, dappled "sunflecks" that illustrate plants' ever-changing light environment. Crops harness light energy to fix carbon dioxide into food via photosynthesis. In a special issue of *Plant Journal*, a team from the University of Illinois reports a new mathematical computer model that is used to understand how much yield is lost as soybean crops grapple with minute-by-minute light fluctuations on cloudy and sunny days.

"Soybean is the fourth most important crop in terms of overall production, but it is the top source of vegetable protein globally," said Yu Wang, a postdoctoral researcher at Illinois, who led this work for Realizing Increased Photosynthetic Efficiency (RIPE). "We found that [soybean](#) plants may lose as much as 13 percent of their productivity because they cannot adjust quickly enough to the changes in [light](#) intensity that are standard in any crop field. It may not sound like much, but in terms of the global yield—this is massive."

RIPE is an international research project that aims to improve photosynthesis to equip farmers worldwide with higher-yielding crops needed to ensure everyone has enough food to lead a healthy, productive life. RIPE is sponsored by the Bill & Melinda Gates Foundation, the U.S. Foundation for Food and Agriculture Research (FFAR), and the U.K. Government's Department for International Development (DFID).

Past models have only examined hour-by-hour changes in light intensity. For this study, the team created a dynamic computational ray-tracing [model](#) that was able to predict light levels to the millimeter across every leaf for every minute of the day in a flowering soybean crop. The model also takes into account two critical factors: photoprotection and Rubisco activase.

Photoprotection protects plants from sun damage. Triggered by high

light levels, this process dissipates excess light energy safely as heat. But, when [light levels](#) drop, it can take minutes to hours for photoprotection to relax, or stop—costing the plant potential yield. The team evaluated 41 varieties of soybean to find out the fastest, slowest, and average rate from induction to the relaxation of photoprotection. Less than 30 minutes is considered "short-term," and anything longer is "long-term" photoprotection.

Using this new model, the team simulated a sunny and cloudy day in Champaign, Illinois. On the sunny day, long-term photoprotection was the most significant limitation of photosynthesis. On the cloudy day, photosynthesis was the most limited by short-term photoprotection and Rubisco activase, which is a helper enzyme—triggered by light—that turns on Rubisco to fix carbon into sugar.

The RIPE project has already begun to address photoprotection limitations in soybean and other [crops](#), including cassava, cowpea, and rice. In 2016, the team published a study in *Science* where they increased the levels of three proteins involved in photoprotection to boost the productivity of a model crop by 14-20 percent. In addition, the RIPE team from the Lancaster Environment Centre at Lancaster University is seeking better forms of Rubisco activase in soybean and cowpea. The RIPE project and its sponsors are committed to ensuring Global Access and making these technologies available to the farmers who need them the most.

"Models like these are critical to uncovering barriers—and solutions—to attain this crop's full potential," said RIPE Director Stephen Long, Ikenberry Endowed University Chair Plant Biology and Crop Sciences at Illinois' Carl R. Woese Institute for Genomic Biology. "We've already begun to address these bottlenecks and seen significant gains, but this study shows us that there is still room for improvement."

More information: Johannes Kromdijk et al, Improving photosynthesis and crop productivity by accelerating recovery from photoprotection, *Science* (2016). [DOI: 10.1126/science.aai8878](https://doi.org/10.1126/science.aai8878)

Yu Wang et al. Photosynthesis in the fleeting shadows: An overlooked opportunity for increasing crop productivity?, *The Plant Journal* (2020). [DOI: 10.1111/tpj.14663](https://doi.org/10.1111/tpj.14663)

Provided by Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign

Citation: Boost soybean yields by adapting photosynthesis to fleeting shadows, according to model (2020, February 24) retrieved 26 April 2024 from <https://phys.org/news/2020-02-boost-soybean-yields-photosynthesis-fleeting.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.