


Forest soils release more carbon dioxide than expected in rainy season

February 14 2020, by Francisco Tutella



Current carbon cycle models may underestimate the amount of carbon dioxide released from the soil during rainy seasons in temperate forests like those found in the northeast United States, according to Penn State researchers. Credit: Caitlin Hodges

Current carbon cycle models may underestimate the amount of carbon dioxide released from the soil during rainy seasons in temperate forests like those found in the northeast United States, according to Penn State researchers.

Tree roots and microbes use [oxygen](#) to convert [organic carbon](#) in the [soil](#) to carbon dioxide (CO₂) for energy through a process called [aerobic respiration](#). This release of CO₂ from soils to the atmosphere represents the largest flux of carbon from terrestrial ecosystems, making it a key component of the global carbon budget. Aerobic respiration is the dominant process contributing to this flux, but the researchers found that under wet conditions, [anaerobic respiration](#)—or respiration without oxygen—also significantly contributes to the flux.

"In current models, amounts of carbon dioxide and oxygen are controlled by the consumption of oxygen and the production of carbon dioxide through aerobic respiration," said Caitlin Hodges, a doctoral candidate in the Department of Ecosystem Science and Management. "It's usually a one-to-one relationship of consumption and production. But we found that, especially in summer, there was a significant signal of anaerobic respiration caused by the roots having a higher demand for oxygen and outcompeting the microbes. The microbes then have to switch over to using anaerobic respiration."

One benchmark for interpreting soil processes affected by soil carbon dioxide and oxygen is to calculate the apparent respiratory quotient (ARQ), which combines carbon dioxide and oxygen concentrations into one value.

"If ARQ equals one, that means that aerobic respiration is the controlling process," said Hodges. "If there is a significant deviation from one, then that tells us something else is controlling the soil gas concentrations."

The scientists studied soil respiration in a shale watershed and a sandstone watershed in the National Science Foundation-funded Susquehanna Shale Hills Critical Zone Observatory. They measured soil carbon dioxide and oxygen levels about eight and 16 inches belowground and just above the bedrock layer where the soil ends.

"Caitlin interpreted gas sampled from soil somewhat like a policeman interprets the breathalyzer test of a tipsy driver," said Susan Brantley, distinguished professor of geosciences and director of the Earth and Environmental Systems Institute (EESI) at Penn State. "The chemistry of the gas stuck in the soil yields a picture of what the bacteria are doing."

The team found that ARQ sometimes signaled significant anaerobic respiration by the microbes. During anaerobic respiration, the microbes shifted from using oxygen to oxidized metals, like iron and manganese, for growth.

"When we see large numbers in the ARQ, it means we have more carbon dioxide than the oxygen levels would suggest," said Jason Kaye, professor of soil biogeochemistry. "How can that happen? It can happen because CO₂ is being produced without oxygen consumption, and that's exactly what an anaerobic process is. That's what these large numbers mean. You're seeing more carbon [dioxide](#) than you would expect from aerobic respiration."

The Penn State team's research, reported in the *Soil Science Society of America Journal*, is the first to use ARQ to find evidence of a seasonal pattern of anaerobic respiration in temperate forests.

The researchers also calculated the total amount of [carbon dioxide](#)—36 grams per meter squared—that leaves the soil system every year due to anaerobic respiration. They said the conservative estimate constitutes 10% of all respiration done by soil microbes at their research sites,

which is a large number since scientists don't think of these humid [temperate forests](#) as posting much anaerobic respiration.

"It's expected that the northeastern United States will have increased rainfall with climate change," said Hodges. "We expect that this anaerobic [respiration](#) will become a more dominant process in these forest systems, and our soil [carbon](#) models don't yet take this into account."

Brantley said novel approaches like soil gas sampling are needed to understand how soils will respond to changing climate.

More information: Caitlin Hodges et al. Soil CO and O Concentrations Illuminate the Relative Importance of Weathering and Respiration to Seasonal Soil Gas Fluctuations, *Soil Science Society of America Journal* (2019). [DOI: 10.2136/sssaj2019.02.0049](https://doi.org/10.2136/sssaj2019.02.0049)

Provided by Pennsylvania State University

Citation: Forest soils release more carbon dioxide than expected in rainy season (2020, February 14) retrieved 3 May 2024 from <https://phys.org/news/2020-02-forest-soils-carbon-dioxide-rainy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
