

# **GPS for chromosomes: Reorganization of the genome during development**

February 28 2020



Embryonic development of the nematode *C. elegans*. Credit: University of Basel, Biozentrum

The spatial arrangement of genetic material within the cell nucleus plays an important role in the development of an organism. A research team from the University of Basel, in collaboration with scientists from Harvard University, has developed a method to trace the chromosomes in individual cells. Using this method, they have now been able to demonstrate that chromosomes reorganize during embryonic



development. The study has recently been published in Molecular Cell.

Our body is made up of a wide variety of cells with the most diverse functions. Irrespective of being heart, liver or nerve cells, however, they all contain the same genetic information. The reason why cells develop differently is that only parts of their <u>chromosomes</u> are read. This results in some genes being active while others, in contrast, are silent.

For gene activation, both the way the genes are packaged as well as their spatial organization in the <u>cell nucleus</u> play a decisive role. Prof. Susan Mango's team at the Biozentrum of the University of Basel has now investigated this 3-D architecture more closely. Using a novel technique, they were able to trace individual chromosomes during <u>embryonic</u> <u>development</u> in nematodes and show that they rearrange themselves during the early phase.

#### **Chromosome arrangement is not random**

If stretched out, all the DNA molecules of a cell would reach about two meters in length. So the DNA must be densely packed to fit into a cell nucleus of only few micrometers in size. The DNA strands are very tightly coiled and twisted to form space-saving structures, called chromosomes. The packaging and the arrangement of the DNA of the chromosomes determines the activity of genes.

In their study, the researchers led by Prof. Susan Mango traced individual chromosomes and investigated their organization during early embryonic development. Embryonic cells of the nematode *C. elegans* served as a model. "Using a novel technique, we were able to follow the spatial rearrangement of chromosomes in single cells at the beginning of embryogenesis," says Mango. "The advantage of this method is that the <u>cells</u> and tissue remain completely intact."



## Early chromosomes resemble a barbell

It is well know that chromosome regions with similar functional properties contact each other and interact. This means that chromosome domains segregated into two compartments, active and inactive. "During early embryogenesis, however, the chromosomes are organized differently," says Ahilya Sawh, first author of the study. "In the early embryo, they are organized into an unconventional barbell-like structure, with inactive compartments separated by a central active region." The researchers discovered that the nuclear lamina—a protein mesh lining the inner surface of the cell nucleus—is required to achieve this barbell arrangement. The lamina is attached to the inactive sections and stretches the chromosome.

### **Chromosomes reorganize during embryogenesis**

"Only at a later stage of embryonic development, when the germ layers develop, we actually see the well-known segregation into an active and inactive region," explains Mango. "Using chromosome tracing, we were able to map the whole 3-D chromosome architecture and could show for the first time that chromosomes rearrange during early development, a maturation process that requires the nuclear lamina."

The reorganization of the chromosomes accompanies cell maturation and represents a milestone in the development of a complex organism. The correct chromosomal architecture is crucial to prevent developmental disorders.

**More information:** Ahilya N. Sawh et al, Lamina-Dependent Stretching and Unconventional Chromosome Compartments in Early C. elegans Embryos, *Molecular Cell* (2020). <u>DOI:</u> <u>10.1016/j.molcel.2020.02.006</u>



#### Provided by University of Basel

Citation: GPS for chromosomes: Reorganization of the genome during development (2020, February 28) retrieved 1 May 2024 from <u>https://phys.org/news/2020-02-gps-chromosomes-genome.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.