


Ionotronic technology that doesn't require liquid electrolytes

February 14 2020, by Bob Yirka



Formation of an IDL at the interface of two oppositely charged ionoelastomers. (A) Schematic illustration of a polyanion/polycation junction. High–surface area carbon nanotube electrodes are embedded within each ionoelastomer, resulting in low-impedance (high-capacitance) EDLs. (B) Chemical structures of the polyanion ES and polycation AT. (C) Nyquist plot and (D) Bode phase plot of acimpedance measurements. The inset in (C) shows an enlargement of the Nyquist



plot in the low-impedance region. Gray lines represent fits of the equivalent circuit model shown in the inset of (D) to the ac-impedance data, where RC, RB, and CB correspond to contact resistance, bulk resistance, and bulk polarization capacitance, respectively. A constant phase element (CPE) is used to describe the EDL (for ES/ES and AT/AT) or the IDL (for ES/AT). Z', real part of complex impedance; Z'', imaginary part of complex impedance. Credit: *Science* (2020). DOI: 10.1126/science.aay8467

A team of researchers from the University of Massachusetts and Harvard University has developed a type of ionotronic technology that does not require liquid electrolytes. In their paper published in the journal *Science*, the group describes several ionic devices they built and possible uses for them. Dace Gao and Pooi See Lee with Nanyang Technological University have published a Perspective piece outlining the work by the team in the same journal issue.

In recent years, companies that make <u>consumer electronics devices</u> have been promising products that users can bend—this would allow for the development of products such as wearable devices. Thus far, there have been very few <u>new products</u>, however. Because of that, scientists have been turning their attention to ionotronic technology in which the <u>charge</u> <u>carriers</u> are ions instead of electrons. But it, too, has met with some roadblocks, most notoriously, the need for liquid electrolytes, which can leak or suffer losses due to evaporation. In this new effort, the researchers in Massachusetts have found a way to create ionotronic devices that do not require liquid electrolytes, perhaps paving the way for new kinds of consumer electronic devices.

To create their new devices, the researchers used two stretchy, rubbery ionoelastomers —polycations and polyanions. With the polycation, positively charged sulfate groups were fixed to the elastomer chains while the negative counterions could easily flow. The polyanions were



made in the same way but with the charges reversed. By combining them, the researchers were able to create a <u>device</u> that allowed current to move in just one direction—a diode analog. They also created a transistor by placing the polycation between two polyanions. And they built a transducer by placing a polyanion atop a polycation and then adding a VHB dielectric layer on top and a carbon nanotube on the bottom.

All of the devices built by the team were both soft and stretchable suggesting that devices with similar materials could soon start showing up in consumer devices. Gao and See Lee suggest that the technology could also be used to create an interface between a machine and a human, noting that both use ions as signal carriers.

More information: Hyeong Jun Kim et al. Ionoelastomer junctions between polymer networks of fixed anions and cations, *Science* (2020). DOI: 10.1126/science.aay8467

© 2020 Science X Network

Citation: Ionotronic technology that doesn't require liquid electrolytes (2020, February 14) retrieved 27 April 2024 from <u>https://phys.org/news/2020-02-ionotronic-technology-doesnt-require-liquid.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.