


# **Researchers discover new mechanism for the coexistence of species**

February 19 2020



Cartoon of experiment showing coexistence between bacteria (red and blue) that cyclically colonize nutrient patches (green). Credit: AMOLF


Researchers from the AMOLF institute in Amsterdam and Harvard have shown that the ability of organisms to move around plays a role in stabilizing ecosystems. In their paper published 19 February 2020 in *Nature*, they describe how the competition between 'movers' and 'growers' leads to a balance in which both types of bacteria can continue to exist alongside each other.

We are all too familiar with the threats to our planet's ecosystems: global warming, <u>forest fires</u>, nitrogen deposition, biodiversity decline, and even mass extinctions. But what actually makes ecosystems stable or fragile? What prevents one species from outcompeting all others, and hence driving them to extinction?



These questions have captivated biologists since Darwin. We have learned that food webs and cooperation between species are key pieces to this puzzle, because they help explain how species depend on one another to survive. Now, a group of biophysicists from the Netherlands and the U.S. have advanced a startling finding: The active movement of organisms can also drive ecosystem diversity and stability through a remarkably simple mechanism that does not require food webs or cooperation.

"Movement is fundamental to essentially all organisms—even plants move by <u>seed dispersal</u>," says Sander Tans at the AMOLF institute in Amsterdam. "Bacteria are well known to move actively. Our experiments show how this movement can keep different bacterial species, typically called strains, together in a larger population. There is a rich literature on the possible roles of movement in such coexistence of species, but direct experiments that can exclude other explanations were lacking.



Bacterial strains A and B were show to both outcompete the other when rare, and hence coexist. Credit: AMOLF




## The coexistence paradox—what prevents extinctions in a competitive world?

To find bacterial species that might form a minimal stable ecosystem in the lab, the Ph.D. student Sebastian Gude took two species (also called strains) from the gut of the same animal. If both survived there, perhaps they would also do so in his experiments. To follow their competition, he colored one blue and the other red. However, all his first attempts failed in the beginning. One of the two strains always lost the competition when grown together, with these <u>bacteria</u> always producing fewer offspring than the other. This blue 'loser' strain would thus become outnumbered by the red 'winner' strain, and was ultimately driven to extinction.

Gude's luck took a dramatic upturn, however, when he changed the design of the experiment. He took the sugary liquid in which the bacteria normally grew, and turned it into a gel reminiscent of a jelly desert. When the blue 'loser' bacteria growing on the gel became outnumbered by the reds, they started to produce more offspring than the reds, and were thus winning. But in turn, the reds also became more competitive when rare. In this way, both strains escaped extinction, and hence coexisted together. These results underscore the fundamental paradox of the coexistence debate: What causes losers that are close to extinction to suddenly start winning?






Fluorescence microscopy of bacterial populations across the gel after the competition. Credit: AMOLF

#### **Getting territorial**

To resolve this conundrum, Gude followed the competition by making movies. "The results were quite striking," says Tom Shimizu. "We saw the populations migrating into the gel like a wave, where they multiplied using the sugars they encountered. Initially, red dominated the expansion and blue was barely seen. But then the red advance suddenly stopped just when the blues emerged, and were seen to overtake the reds' front, where they formed a thin layer. After that, the wave was blue only. The blue bacteria could thus proliferate alone in the deeper regions of the gel, freed from competition from the reds that could not reach that far. This also explained the coexistence: Whenever blue were rare, they could build up their population in the deeper regions of the gel."





Accelerated movie of competing bacterial strains. Credit: AMOLF


### Should I spread or should I grow?

But how did the blue bacteria organize themselves and efficiently confine the red? Did they signal to each other or secrete toxins, as is known for some bacteria? In trying to address these questions the team discovered a rather different mechanism. The blue bacteria were indeed worse at proliferating, as they lost in a direct competition. But they compensated by migrating faster. By reaching the farther regions first, they could finish the local sugars. Hence, they gave the red no chance there, and could thus block their advance, a bit like in a scorched earth strategy.

Tans: "Some bacteria apparently are good at proliferating, and others at migrating. But they cannot excel at both. This makes sense because both



activities cost a lot of energy. Such specialization is often observed, though effects on coexistence are often difficult to prove. Here we could manipulate the capacity to migrate and proliferate by genetic engineering, and show it alone is enough for coexistence. Other mechanisms like exchanging toxins or dependencies like in <u>food webs</u> are thus not required per se."



Slowed-down movie of motile E. coli cells. Credit: AMOLF

#### **Disturbance ecology**

Of course, in the real world, bacteria cannot count on encountering bowls of Jello left out of the fridge. Luckily for them, they don't need to, as pristine pastures of fresh nutrients do arise more often than expected. Much like the vegetation that colonize cleared soil after forest fires, many bacteria grow on patches of resources, from fallen fruits to decomposing animals, or—for those microbes that inhabit your gut—the lunch you just had.

Shimizu says, "We demonstrated this migration-proliferation mechanism in motile bacteria. But what we found is for instance evocative of models of plant ecology where fast-growing plants compete with plants that invest more in spreading their seeds." He added that there is a lot to be



learned about more complex scenarios. "There is a lot of interest in explaining the diversity of bacteria in the human gut, which we now know, thanks to modern DNA analysis, is intimately related to our health. Our findings suggest that looking at motility genes and their spatial distribution within the gut might help explain some of that diversity."

**More information:** Bacterial coexistence driven by motility and spatial competition, Sebastian Gude, Erçağ Pinçe, Katja M. Taute, Anne-Bart Seinen, Thomas S. Shimizu, and Sander J. Tans, *Nature* 2020, <u>DOI:</u> 10.1038/s41586-020-2033-2

Provided by AMOLF

Citation: Researchers discover new mechanism for the coexistence of species (2020, February 19) retrieved 2 May 2024 from <u>https://phys.org/news/2020-02-mechanism-coexistence-species.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.